

# Middle Potomac River Watershed Assessment: Potomac River Sustainable Flow and Water Resources Analysis



April 2014



# Middle Potomac River Watershed Assessment: Potomac River Sustainable Flow and Water Resources Analysis

United States Army Corps of Engineers, Baltimore District

The Nature Conservancy

Interstate Commission on the Potomac River Basin

Final Report

April 2014

Recommended citation:

U.S. Army Corps of Engineers, The Nature Conservancy, and Interstate Commission on the Potomac River Basin. 2014. Middle Potomac River Watershed Assessment: Potomac River Sustainable Flow and Water Resources Analysis. Final Report. 107 pp. and 11 appendices.

Cover photograph: Great Falls on the Potomac River as seen from Virginia. Jim Palmer (ICPRB)

# Abstract

---

The Middle Potomac River Watershed Assessment (MPRWA) was a collaborative effort to assess the relationship between streamflow alteration and ecological response in the Potomac River and its tributaries in a study area defined as the Middle Potomac. The assessment is comprised of five distinct components: (1) a large river environmental flow needs assessment, (2) a stream and small rivers environmental flow needs assessment, (3) a projection of future water uses, (4) a stakeholder engagement process, and (5) development of a concept or scope for a strategic comprehensive plan for watershed management. This information can be used to balance and mitigate water use conflicts and prevent ecological degradation.

To assess flow needs in the Potomac watershed, the Ecologically Sustainable Water Management (ESWM) approach was used for large rivers and the Ecological Limits of Hydrologic Alteration (ELOHA) framework was adapted for streams and small rivers.

In the large rivers included in this study, based on currently available information, there has been no discernible adverse ecological impact on focal species due to human modification of flows. As a precautionary measure, the team recommended that the current large river flow regime be maintained for the entire range of flows as defined by 20 flow statistics based on a 21-year period of record (1984-2005).

In streams and small rivers, strong relationships were found between urbanization (impervious surface), hydrologic alteration, and ecological impacts. Land use was found to be a more significant cause of hydrologic alteration than water withdrawals and impoundments at the present time. Impervious surfaces move rainfall and snowmelt more rapidly into rivers and streams than surfaces with natural vegetation, causing increased flashiness (rate of change). Increased flashiness correlates with declines in the status of stream macroinvertebrate, the streamflow assessment's indicator taxa. These taxa were not found to be sensitive to changes in low magnitude flows. Analyses of other taxa more sensitive to low magnitude flows, such as fish and mussels, could provide a more complete picture of biological response to flow alteration for water use decisions. Basin-wide datasets of these groups are not collected at present.

The impact on flow of future changes in water use was assessed through six scenarios: a) three different forecasts of per capita domestic water use; b) climate change; c) hot and dry summer conditions; and d) conversion of power plants to closed cycle operation. There was no regional pattern of flow alteration that applied to all scenarios and, within scenarios, impacts on flow varied for each subwatershed's unique combination of land and water uses.

Stakeholder education and engagement are key elements in the process of transferring scientific findings to management and policy. Workshops, webinars, and technical meetings were convened in order to obtain advice on methods, interpret draft results, and share findings with government agencies at all levels and other parties with significant roles in water management. This dialogue and information sharing contributes to state water management decision making processes and to the ongoing dialogue among stakeholders about ensuring the sustainable use of stream flows for all purposes.

Water use and basin development in each of the Potomac watershed's jurisdictions affects the other jurisdictions. The final component of this study was to scope a basin-wide comprehensive plan process. Because there is no single agency with regulatory authority for water planning across the Potomac watershed, the concept for a future comprehensive plan is based on a collaborative, stakeholder driven model.

## Table of Contents

Abstract .....	i
List of Tables .....	iv
List of Figures.....	iv
Executive Summary .....	ES-1
Chapter 1. Introduction .....	1
Chapter 2. Existing Conditions.....	6
2.1. Physiography .....	6
2.2. Climate .....	7
2.3. Hydrology .....	7
2.4. Impoundments.....	8
2.5. Water Withdrawals and Discharges .....	10
2.6. Land Uses .....	10
2.7 Ecological Community .....	11
Chapter 3. Large River Environmental Flow Needs .....	14
3.1. Introduction .....	14
3.2. Large River Flow Needs Assessment Approach .....	15
3.3. Findings of the Large River Flow Needs Assessment .....	18
3.4. Conclusions for Large River Flow Needs Assessment .....	20
3.5. Large River Information Gaps and Monitoring Needs .....	21
Chapter 4. Future Water Use Projections .....	23
4.1. Methods for Future Water Use Projections .....	23
4.2. Findings for Future Water Use Projections.....	25
4.3. Comparison to Previous Studies .....	27
4.4 Summary .....	27
Chapter 5. Stream Environmental Flow Needs .....	29
5.1. Ecological Limits of Hydrologic Alteration (ELOHA) .....	30
5.2. Findings for Stream Environmental Flow Needs: Interpretation and Use of Relationships.....	33
5.2.1. Magnitude.....	34
5.2.2. Duration .....	34
5.2.3. Frequency .....	37
5.2.4. Rate of Change .....	37
5.2.5. Possible Mechanisms Underlying FA-E Relationships .....	39
5.2.6. Other Environmental Stressors .....	41
5.2.7. Using the FA-E Curves.....	43
5.3. Discussion and Conclusions for Stream Environmental Flow Needs .....	46
5.3.1. Land Use Change Thresholds of Potential Concern .....	46
5.3.2. Biological Thresholds of Potential Concern .....	48
5.4. Technical Information Gaps, Research, and Analysis Needs .....	51

## Middle Potomac River Watershed Assessment

Chapter 6. Stakeholder Outreach and Implementation Options.....	53
6.1. Moving the Science into Management .....	53
6.2. Stakeholder Participation.....	54
6.3. Computer-Based Evaluation Tool.....	55
6.4. Potomac Basin Comprehensive Water Resources Plan.....	56
6.5. Discussion.....	57
Chapter 7. Conclusions .....	58
7.1. Introduction .....	58
7.2. Study Findings.....	58
7.2.1. The Potomac Large River Environmental Flow Needs Assessment.....	59
7.2.2. Future Water Use Projections .....	59
7.2.3. Stream and Small Rivers Environmental Flow Needs Assessment.....	59
7.2.4. Stakeholder Engagement Process.....	62
7.2.5 Summary.....	62
7.3. Study Products .....	63
7.4. Lessons Learned .....	66
7.5. Information Needs.....	67
7.5.1. Research and Analysis .....	67
7.5.2. Monitoring .....	68
7.5.3. Modeling.....	69
7.6. Applicability to Jurisdictions.....	69
7.7. Next Steps.....	70
Supplemental Tables.....	72
Literature Cited.....	76
Glossary.....	86
Abbreviations and Acronyms.....	88
Team Members and Contributors .....	90

## Appendices

A	Large River Environmental Flow Needs Report
B	Water Withdrawals and Consumptive Use in the Potomac River Report
C	Compilation of Measured Stream Data
D	Application of ELOHA
E	Development and Refinement of Hydrologic Model
F	Flow Metric Testing
G	Stream Classification
H	Basin-Wide Hydrologic Alteration Assessment
I	Development of Stream Flow Alteration – Ecological Relationships
J	Stakeholder Engagement
K	Implementation Options

**List of Tables**

Table 1. Organization of the MPRWA report..... 5

Table 2. Impoundments in the Middle Potomac River study area and North Branch Potomac River holding more than 10 percent capacity of annual flow volume. .... 9

Table 3. Withdrawals and consumptive use by sector in the Middle Potomac River study area. .... 26

Table 4. Apparent thresholds of potential concern for different land use combinations in the Middle Potomac simulated results. .... 49

Table 5. Summary of flow alteration – macroinvertebrate relationships identified in this study. .... 60

Table 6. Summary of major products. .... 64-65

Supplemental Tables

Table A. Flow metrics ..... 72-73

Table B. Macroinvertebrate family-level metrics. .... 74-75

**List of Figures**

Figure 1. Middle Potomac watershed study area. .... 2

Figure 2. Potomac River basin boundaries. .... 6

Figure 3. Land use in the Potomac River basin. .... 11

Figure 4. Conceptual diagram of the major steps in developing Flow Alteration-Ecology (FA-E) relationships for the Middle Potomac study area. .... 30

Figure 5. Conceptual diagram of biological community health. .... 32

Figure 6. FA-E relationships for positive alteration (increase) in annual 3-day maximum flow. .... 35

Figure 7. FA-E relationships for negative alteration (decrease) in the high flow duration metrics DH17 (A) and MH21 (B). .... 36

Figure 8. FA-E relationships for negative alteration (decrease) in low pulse duration. .... 36

Figure 9. FA-E relationships for positive alteration (increase) in high pulse count. .... 38

Figure 10. FA-E relationships for positive alteration (increase) in flashiness. .... 38

Figure 11. Chessie BIBI vs percent alteration in flashiness. .... 42

Figure 12. An illustration of how watershed characteristics can be linked to flow alteration and flow alteration linked to biological response. .... 44-45

Figure 13. Simplified flow chart of the steps used to select the final sample locations, flow metrics, and macroinvertebrate metrics. .... 47

# Executive Summary

The Middle Potomac River Watershed Assessment (MPRWA) was a collaborative effort to assess streamflows in the Potomac River and its tributaries in a study area defined as the Middle Potomac (Figure ES-1). Beginning in May 2009, and working with Potomac River watershed experts, the U.S. Army Corps of Engineers (USACE), the Interstate Commission on the Potomac River Basin (ICPRB), and The Nature Conservancy (TNC) collaborated to identify key ecological needs related to streamflow, existing and future impacts of human activities on flow, and the potential effects of climate change on the watershed’s hydrology. As technical partner, ICPRB coordinated and conducted the study’s scientific analyses.

To assess flow needs in the Potomac watershed, ICPRB adapted the Ecologically Sustainable Water Management (ESWM) approach (Richter et al. 2003) for large rivers and the Ecological Limits of Hydrologic Alteration (ELOHA) framework (Poff et al. 2010) for streams and small rivers. Expert workshop participants and a technical advisory group, representing watershed jurisdictions and interested Federal agencies, provided input and feedback to inform the analyses. In addition to analyses and findings, the report presents research, monitoring, and modeling needs to complement the assessment, and provides lessons learned to inform other environmental flows analyses.

For large rivers, the project team concluded, based on currently available information, there has been no discernible adverse ecological impact on focal species due to human modification of flows. As a precautionary measure, the team recommended that the current large river flow regime be maintained for the entire range of flows as defined by 20 flow statistics based on a 21-year period of record (1984-2005).

In streams and small rivers, strong relationships were found between urbanization (impervious surface), hydrologic alteration, and ecological impacts. Land use was found to be a more significant cause of hydrologic alteration than water withdrawals and impoundments. Impervious surfaces, in contrast to natural soils and vegetation, move rainfall and snowmelt more rapidly into rivers and streams, causing increased flashiness (rate of change). Increased flashiness correlates with declines in stream macroinvertebrate status, the streamflow assessment’s indicator taxa. These taxa, however, were not found to be sensitive to changes in low flows. Analyses of other taxa more sensitive to low flows, such as fish and mussels, could provide a more complete picture of biological response to flow alteration for water use decisions, but basin-wide datasets of these groups are not collected.




Figure ES-1. Middle Potomac watershed study area.

## Project Background

The purpose of the MPRWA was to investigate the environmental flow<sup>1</sup> needs of rivers and streams in the Middle Potomac watershed, the area defined as the North Branch confluence to Occoquan Bay (Figure ES-1). Changes in natural flows are caused by water withdrawals, discharges, impoundments, land use change, and climate change. Environmental flows may differ from natural flows in order to accommodate human uses of water. Maintaining or restoring an environmental flow regime – including the magnitude, duration, timing, frequency, and rate of change of hydrologic events such as high and low flows – is important for stream health.

Many river ecologists consider flow to be a master variable for aquatic ecosystems because it influences water quality, biotic interactions, and the availability of food and habitat for fish and other species. However, flow is only one of multiple factors affecting biological community health, and flow itself is influenced by a wide range of natural and human-influenced watershed conditions. One challenge of this analysis was to distinguish biological changes caused by flow alteration from changes caused by other factors, such as poor water quality and instream habitat disturbance.

The Potomac River is the second largest tributary to the Chesapeake Bay. Its mainstem flows southeast, cutting through the Appalachian Mountains and Piedmont, becoming tidal as it enters the Atlantic Coastal Plain. Its watershed is shared by four states (Maryland, Pennsylvania, Virginia, and West Virginia) and the District of Columbia. Few dams regulate flow in the streams and large rivers of the Potomac River basin compared to other eastern U.S. river systems. Most of the 481 impoundments identified in the watershed are run-of-river facilities and only minimally alter flow patterns. For this reason, the Potomac River presents a rare opportunity to be proactive in defining the environmental flows required to sustain natural diversity and ecosystem functions while meeting the needs of a growing regional human population. Growth is expected to convert forest and farmland into hardened landscapes, increase levels of runoff and pollution to the river and the Bay, and increase electricity demands and consumptive uses of water. This study is timely considering the watershed jurisdictions' development of state water management plans and policies and the potential for more droughts and catastrophic floods due to global climate change.

## Methods and Findings

The MPRWA was comprised of five distinct components: (1) a large river environmental flow needs assessment, (2) a projection of future water uses, (3) a stream and small rivers environmental flow needs assessment, (4) a stakeholder engagement process, and (5) development of a concept and process for a basin-wide comprehensive plan for watershed management. This assessment generated datasets and analyses that can help the Potomac River watershed jurisdictions protect environmental flows.

### Large river environmental flow needs assessment

The large river assessment adapted the ESWM approach (Richter et al. 2003), which uses available hydrologic data, hydro-ecological literature, and expert judgment to determine environmental flows. A hydrologic risk assessment was conducted to identify large river segments and tributaries at greatest risk of hydrological alteration from several risk factors, including geology and current and projected future water and land use. The sub-basins deemed most at risk of hydrologic alteration

---

<sup>1</sup> Environmental flows are defined as the seasonally variable flows of water that sustain healthy river ecosystems and the goods and services that people derive from them.

were the Occoquan, Monocacy, Aquia, Opequon, Mattawoman, Saint Mary's, Goose, Conococheague, Antietam, North Branch, Cameron Run, and Accotink. The mainstem river segments at greatest risk extended from the Potomac River above Paw Paw to the Potomac River above Little Falls.


The team chose to focus the large river flow needs assessment on four large river segments and two large tributaries: (1) Potomac River mainstem from the Shenandoah River confluence to Point of Rocks, (2) Potomac River mainstem from Point of Rocks to Great Falls, (3) Potomac River mainstem from Great Falls to Chain Bridge, (4) tidal freshwater Potomac River estuary from Chain Bridge to Occoquan Bay, (5) Monocacy River mainstem, and (6) Opequon Creek mainstem. These reaches are at highest risk for hydrologic alteration and were selected as regions of special interest for varying reasons including current development pressures, land use, physiography, and other criteria. Environmental flows for non-tidal, free-flowing rivers were based on fish, mussels, and instream and riparian plant needs; flows for the tidal freshwater estuary were based on fish, benthic macroinvertebrates, zooplankton, and phytoplankton needs.

Based on a review of available hydro-ecological literature, hydrologic data, and expert opinion, the project team found no current discernible adverse impact on key biotic taxa in large rivers due to human modification of flow. Potomac River flows, however, are naturally highly variable due to hydro-meteorological factors, so it is difficult to separate short-term, weather-related impacts from long-term, anthropogenic impacts on biota. As a precautionary measure, the team recommended that the current large river flow regime be maintained for the entire range of flows. The team also recommended maintaining the existing 100-mgd minimum flow requirement for Little Falls and 300-mgd minimum flow recommendation for Great Falls. Participants at a September 2010 workshop reviewed the draft assessment and recommended that a technical workgroup be convened to design a monitoring program to fill information gaps for these systems.

### **Projection of future water uses**

The future water use component of this study estimated the magnitude of future increases in water withdrawals and especially consumptive uses, identified the principal causes of those increases, and examined how those increases might affect the stream-flow regime through 2030. Water withdrawal is water diverted or removed from a source, a large percentage of which may be returned to the source, such as through wastewater treatment plant discharges. The portion of water withdrawal that is removed and is not available for downstream use by humans or the aquatic ecosystem is termed "consumptive use." Consumptive use is water withdrawn and permanently removed from surface or groundwater sources because it has been evaporated, transpired by plants, incorporated into products or crops, consumed by humans or livestock, transferred out of the watershed, or otherwise removed from the watershed (Kenney et al. 2009).

The rate of growth in withdrawals and consumptive use is strongly dependent on assumptions about individual behavior (per capita rates of consumption) and about future changes in the industry, power, and agriculture sectors. Currently, domestic and public supply (water withdrawn by public and private water suppliers and delivered to users) in the MPRWA study area (includes the Washington, D.C. metropolitan area) accounts for 27 percent of the 2,252-mgd total water withdrawals by all sectors, but 84 percent of the 394-mgd total consumptive water use by all sectors (Figure ES-2). For non-drought years and assuming no change in the per capita rate of domestic consumption, total consumptive use is predicted to increase to 488 mgd by 2030, a 24-percent increase from 2005. Domestic and public supply would still account for 84 percent of the total



**Figure ES-2. Percentage of surface withdrawals and consumptive use by sector in 2005 for the Middle Potomac River Watershed Assessment study area.**

consumptive use. Scenarios that assume an increase in the per capita domestic water use show higher increases in consumptive use. Water used for power plant cooling currently accounts for 68 percent of surface withdrawals by all sectors but most of that water is returned to streams, and so the power sector accounts for only 12 percent of total consumptive use. Conversion of power plants to closed-cycle cooling is predicted to result in an increase in consumptive use by this sector. Agriculture currently accounts for less than 1 percent of surface withdrawals and 1.3 percent of consumptive use. Consumptive use by agriculture during an extreme drought is predicted to increase by nearly three times, but still is only a few percent of the total basin-wide consumption. A comparison of consumptive use forecasts made by this project with those made by Steiner et al. (2000) is difficult due to differences in data and methods and inherent uncertainties in data sources, but the predicted rates of growth in consumptive use are similar – about one mgd each year.

The Chesapeake Bay Program’s watershed model was adapted and linked with a river transport model to predict current and future daily water flows at ungaged outlets throughout the Potomac River watershed. Results provided a basis to estimate future flow alteration in subwatersheds (equivalent to watershed model river segments) and to make a spatial assessment of watersheds at greatest risk of future hydrologic alteration. Future flow alteration predicted by five modeled scenarios varied across the basin, depending on the assumptions of each scenario, on changes in individual watershed characteristics (land and water use), and on what aspect of the flow regime (as measured by a specific flow statistic) was being considered.

### **Stream and small rivers environmental flow needs assessment**

For the streams and small rivers flow assessment, the ELOHA framework (Poff et al. 2010) was adapted and used to develop flow alteration-ecological response (FA-E) relationships from modeled flow data and a basin-wide biological dataset of benthic macroinvertebrates. This component of the MPRWA study is one of the first attempts to perform a large-scale quantitative analysis using the ELOHA approach. Stream macroinvertebrates were selected as biological response variables because of the larger size, better integration, and broader coverage of macroinvertebrate data across the basin jurisdictions and the lack of comparable fish data at the basin-wide scale.

Macroinvertebrates are a major food source for fish and birds and are commonly used as indicators of stream health.

The team simulated streamflow and flow alteration at biological sampling sites because there are few gages recording flow data in streams where macroinvertebrates are sampled. The hydrologic modeling was done with the Chesapeake Bay Program’s watershed model and the Virginia Department of Environmental Quality’s Online Object Oriented Meta-Model (WOOOMM) routing module. Current and baseline flow time series were simulated for 747 macroinvertebrate sampling locations. The baseline flow time series is defined as modeled current conditions with the influence of water withdrawals, impoundments, and biologically significant hydrologic alterations due to land use removed via modeling. Flow metrics calculated from the simulated time series were tested against observed data. Through a series of analyses six flow metrics and seven biometrics were selected to generate FA-E relationships (Table ES-1). The selected flow metrics represent different aspects of the flow regime, show relationships with watershed and water use factors, and show relationships between flow alteration and biological status.

Watershed factors that alter flow in the Middle Potomac study area were related most often to urbanization, particularly impervious surface area. Flashiness increases sharply from baseline, or relatively unaltered conditions, when the area of impervious surface in a watershed exceeds a threshold of about one percent. Water withdrawals and discharges directly alter flows, but significant impacts on flow metrics were seen only in those watersheds where net withdrawals or net discharges are a large fraction of mean annual flow. Impacts of agriculture on streamflow were not readily apparent in the Middle Potomac study area, but they can confound some hydrologic impacts of impervious surface area.

The FA-E relationships developed through this analysis indicate that for a variety of flow metrics, hydrologic alteration is associated with a decrease in the likelihood of many macroinvertebrate metrics achieving comparatively healthy status (“fair” or better). Higher high flows, more frequent high and low flow events, shorter high and low flow periods, faster rise and fall rates, and more reversals had large and negative impacts on all metrics of macroinvertebrate health. None of the biological metrics responded to change in middle and low magnitude flow metrics (Table ES-2).

The Chessie BIBI, a basin-wide, multi-metric Benthic Index of Biological Integrity developed for stream macroinvertebrates in the Chesapeake Bay watershed, reflects biological condition. Figure ES-3 illustrates that as percent alteration of flashiness increases, biological condition decreases. The black line in the figure is a linear regression at the 90th percentile of the data. It is an approximate indication of the best possible biological status that can be achieved at different levels of flow alteration, given the existing mix of environmental factors affecting stream macroinvertebrates.


**Table ES-1. Macroinvertebrate and flow metrics used to establish FA-E relationships for the Middle Potomac River Watershed Assessment study area.**

Macroinvertebrate metrics	Flow metrics
Chessie BIBI (a multi-metric community index)	3-day maximum
%EPT (pollution-sensitive group)	High flow index MH21
FBI (Hilsenhoff family-level biotic index)	High flow duration DH17
%Chironomidae (pollution-tolerant group)	Low pulse duration
%Scraper (feeding group)	High pulse count
%Clinger (habit group)	Flashiness
SW (Shannon-Wiener diversity index)	

**Table ES-2. Summary of flow alteration – macroinvertebrate relationships identified in the study.**

Alteration in these aspects of the flow regime is associated with →	Degradation in:	Possible mechanisms that could explain the association are:
<ul style="list-style-type: none"> <li>• Higher maximum flows</li> <li>• Shorter duration of high flows</li> <li>• Shorter duration of low flows</li> <li>• More low flow pulses</li> <li>• More high flow pulses</li> <li>• Faster rates of change in flow (flashier)</li> </ul>	All family-level macroinvertebrate metrics tested and the Chessie BIBI multi-metric index	<ul style="list-style-type: none"> <li>• Scour of periphyton and organic matter (food) during high flows</li> <li>• Catastrophic accidental drift during floods</li> <li>• Displacement from habitat and stranding when waters recede</li> <li>• Physical alteration of stream bed habitat</li> <li>• Indirect effects of poor runoff water quality (sedimentation, pollutants)</li> <li>• Interruption of development or dispersal cues</li> </ul>
<ul style="list-style-type: none"> <li>• Lower middle and low magnitude flows, includes median flow, August median flow, summer Q85 flow, baseflow index, 3-day and 1-day annual minima, and 7Q10</li> </ul>	None of the biometrics	<ul style="list-style-type: none"> <li>• Swift recovery due to adaptations to low flow (drought resistant or diapausing life stages)</li> <li>• Multi-voltine (short) life cycles</li> <li>• High mobility, able to find refugia and later recolonize</li> </ul>


As other environmental factors become more stressful (e.g., poor water quality, physical disruption of stream habitat), the “cloud” of data points in Figure ES-3 shifts lower on the plot. Figure ES-4 illustrates how reduced water quality or habitat condition reduces the limit of best possible biological status (represented by the regression line). The regression line in a severely degraded stream (dashed) is lower than the line in an otherwise excellent stream environment (solid). Point (A) indicates where increasing flow alteration lowers biological status below what could be deemed an “acceptable” threshold in an excellent quality stream. Point (B) indicates the threshold is crossed much sooner in an already degraded stream. To accurately forecast change in the biological community caused by a specific change in flow, projections must account for all the natural and anthropogenic factors in addition to flow that impact biological communities. Water quality and habitat improvements can potentially ameliorate the impacts of future flow alteration and the regression line would shift up. Conversely, future flow alteration will further degrade biological status even if water quality and habitat conditions are unchanged. This finding is why the study does not put forth any specific flow alteration limits to protect the environmental flow needs of streams and small rivers in the Middle Potomac River study area.



**Figure ES-3. Relationship between percent alteration of flashiness with Chessie BIBI.**  
The black quantile regression line is significant at the 0.01 level.


This study’s FA-E relationships can be used to evaluate the biological impact of alternative scenarios for watershed development or water use if the impacts of other factors are taken into account. Conditional probability plots that relate flow alteration to the probability that biological status will be fair or better can be used to estimate an increased risk to biological health due to flow alteration. Thresholds in the relationships between flow alteration and land use, developed for this project, can inform managers when changes in specific land uses are likely to begin altering stream flows. For example, proposed (future) increases—or decreases—in watershed impervious surface area can be evaluated for their potential impact on flashiness and stream macroinvertebrate condition (Figure ES-5). Differences in the curves indicate different levels of sensitivity to flow alteration in the macroinvertebrate metrics; however, the general pattern is a decline in biological condition with an increase in flashiness. Plots have been developed for the 42 FA-E relationships (7 macroinvertebrate metrics x 6 flow metrics) developed for the MPRWA.

Although the FA-E relationships developed in the study are generally applicable across the non-tidal Potomac watershed, consideration must be given for the confounding influences of other environmental factors and the innate differences in biometric response strength. In watersheds that hydrologic models can not represent well at this time, such as the small watersheds in the broad corridor of karst geology bisecting the project’s study area, the study’s FA-E relationships should not be applied. In these watersheds, resource managers could apply a presumptive flow protection standard (Richter et al. 2011).



**Figure ES-4. Generalized illustration of the influence of water quality and stream habitat conditions on macroinvertebrate responses to flow alteration.**

The solid line represents a high quality stream and the dashed line represents a degraded stream.



**Figure ES-5. Conditional probability plots of FA-E relationships for positive alteration (increase) in flashiness.**

### **Stakeholder engagement process**

This study provided extensive stakeholder education and engagement to inform and review the analysis and to encourage the future transfer of scientific findings to management and policy. Workshops, webinars, and technical meetings were convened to obtain advice on methods, interpret draft results, and share interim findings with government agencies and other parties with significant roles in water management. The intent was to contribute to water management decision making and to the ongoing dialogue among stakeholders about ensuring sustainable water use for all purposes.

### **Development of basin-wide comprehensive plan concept**

Water use and land development in each of the Potomac River watershed's jurisdictions affect the other jurisdictions. Basin-wide collaboration is needed to protect flow regimes, healthy watersheds, and river and stream health across the watershed. To put this environmental flow assessment in a broader watershed management context, and to promote the development of a basin-wide vision for Potomac River management, this project included the scoping of a basin-wide comprehensive plan process. The process is expected to include four phases: (1) Define roles of participating organizations, establish interdisciplinary oversight committee, and define goals of plan; (2) Assess water resources issues and identify problems; (3) Identify and evaluate alternatives, and recommend practical solutions; and (4) Develop the comprehensive water resources plan document. Because there is no single agency with regulatory authority for water or land use planning across the Potomac River basin, the concept for the comprehensive plan is based on a collaborative, stakeholder driven model, which ICPRB will be pursuing in the future with the watershed jurisdictions.

### **Potential applications of study products and findings**

The MPRWA explored how biological communities respond to hydrologic alteration and other changes in watershed conditions. Its datasets, methods, and analysis results can be used by watershed jurisdictions to assist with water resources protection efforts. Maintaining healthy streams and watersheds, as reflected through the status of stream biological communities, is important to meet Potomac River and Chesapeake Bay restoration goals. Healthy stream communities are more resilient to natural, extreme events and to the multitude of anthropogenic stressors that accompany human activities. They are more capable of processing both natural leaf litter and human pollutants. They provide greater support to organisms higher in the food web, many of which are economically and recreationally important. Healthy stream systems are more capable of providing the many goods and services that humans depend upon, from drinking water supply to wastewater assimilation to flood mitigation to recreation.

Defining limits of acceptable hydrologic alteration towards management of ecological health, a final step in the ELOHA process, is a stakeholder driven decision-making process. Limits of acceptable alteration were not defined in this study. The process of defining societal values and needs, determining desired ecological conditions for specific streams and watersheds, and setting limits to flow alteration can best be pursued at a watershed jurisdictional scale, in accordance with state- and local-level priorities, needs, and regulatory mandates. This study's results can provide a scientific foundation to inform that process. Local water resource managers or planners could go through a series of steps to develop flow recommendations based on the FA-E response relationships identified in this analysis. At the local level, the results may be most applicable for land use planning and zoning. At the state level, the results could influence the development or refinement of state stormwater laws to advance Bay restoration goals, or support the development of new flow-related water quality standards.

# Chapter 1. Introduction

---

**Chapter Summary:** This chapter describes the background and scope of the Middle Potomac River Watershed Assessment (MPRWA) project. The project consisted of five related components. One component—an application of the Ecological Limits of Hydrologic Alteration (ELOHA) process to streams and small rivers in the study area—is described at length in Chapter 5 and several appendices of this report. Three components are described in separate technical reports, included as appendices of this report. Their highlights and conclusions are summarized in chapters 3 (large river environmental flow needs), 4 (future water use projections), and 6 (stakeholder outreach and implementation options). Chapter 2 describes existing conditions in the study area. Chapter 7 presents the overall study conclusions, lessons learned, and next-steps.



The Middle Potomac River watershed encompasses 11,550 mi<sup>2</sup>, or 79 percent, of the Potomac River basin (Figure 1), from the North Branch confluence to Occoquan Bay. This area had a population in 2010 of 5.57 million people and includes much of the Washington DC metropolitan area in the east. The Middle Potomac River watershed has a diverse landscape of mountains, valleys, and coastal plains, with urban, rural, and natural areas. The river is the second largest tributary to the Chesapeake Bay and, with four states (Maryland, Pennsylvania, Virginia, West Virginia) plus the District of Columbia, it is the only tributary shared by a majority of the Bay watershed jurisdictions. Although the official geographic scope of this assessment is the Middle Potomac River watershed, the project team recognized the need, for both hydrologic and ecological considerations, to include the North Branch and the estuarine Potomac for some analytical purposes.

The Potomac River basin has experienced rapid population growth in recent years, growing 15.6 percent between 2000 and 2010. With this growth has come increased development, including new and larger water withdrawals (surface and groundwater) and more urbanization. Stream corridors are increasingly experiencing water resource, or flow, related problems including: degradation and loss of wetlands and fish and wildlife habitat; excessive stream bank erosion; exacerbated flooding; excessive nutrients and sediments; and water supply and management conflicts. Clear connections between land uses, water uses, and flow alteration, and between flow alteration and biological degradation, are quantitatively described in the literature, most recently in publications such as Carlisle and Meador 2007, Kennen et al. 2009, Falcone et al. 2009, Carlisle et al. 2010, Cruise et al. 2010, Cuffney et al. 2010, and Poff and Zimmerman 2010. Continued population growth in the watershed is expected to heighten demand for water and electricity, as well as convert forest and farmland into developed and hardened landscapes and increase levels of runoff and pollution to the river and the Chesapeake Bay. This growth will likely amplify the severity of flow alteration and flow-related impacts. The extent and cumulative impact of these problems is of concern to the region's natural resources agencies.

Compared to other large eastern U.S. river systems, the Potomac River is relatively intact with respect to stream flow, with only a few large dams that significantly change flow. For this reason, the Potomac presents a rare opportunity to evaluate increasing human impacts on flow originating from sources that are more diffuse, such as agricultural and urban land uses, surface and groundwater withdrawals, and discharges. The opportunity is timely considering the watershed

## Middle Potomac River Watershed Assessment

jurisdictions' development of state water management plans and policies, the increasing demand for consumptive use of river water (amount of water withdrawn but not returned), and the potential for greater incidences of droughts or catastrophic floods with global climate change. Approaches exist for developing science-based environmental flows and implementing plans and policies to protect them (e.g., Richter et al. 2003, Poff et al. 2010, Richter et al. 2011) and are being tried in the United States (e.g., Kendy et al. 2012).

The Middle Potomac River Watershed Assessment (MPRWA) was conducted under the authority of Section 729 of the Water Resources Development Act (WRDA) of 1986, as amended. In addition to the Section 729 authority, several Congressional committee resolutions provide additional authority for the U.S. Army Corps of Engineers (USACE) to study the Potomac River watershed. After scoping the watershed assessment, The Nature Conservancy (TNC), the non-federal sponsor, and USACE, Baltimore District, entered into a partnership agreement in May 2009. The Interstate Commission for the Potomac River Basin (ICPRB) was contracted to provide technical expertise and conduct much of the analysis in the MPRWA.

The overall goal was to provide a large, diverse foundation of information useful in implementing future protection and restoration of the basin's water resources over the next several decades. The study included data gathering, flow modeling, ecological investigations, and consultation with




Figure 1. Middle Potomac watershed study area.

stakeholders. The study team considered water and land uses that affect the flow regime and investigated environmental flow<sup>2</sup> needs, landscape changes, and watershed resource management in the Middle Potomac River watershed in the District of Columbia, Maryland, Pennsylvania, Virginia, and West Virginia. Also considered was how future changes in human activities (especially withdrawals, impoundments, and land use development) and climate change might impact the flow regime. Five separate but related study components comprised the MPRWA. They produced:

- 1) A large river environmental flow needs assessment for the mainstem Potomac River (below Harpers Ferry to the confluence with the Occoquan River) and mainstem Monocacy River and Opequon Creek;
- 2) A forecast of future water use to 2030 based on state databases of water use for domestic and public supply, literature estimates of water use by industrial and agricultural sectors, and projections of population, land use, industry, and climate change;
- 3) A stream and wadeable rivers environmental flow needs assessment using the Ecological Limits of Hydrologic Alteration (ELOHA) approach and based on observed benthic macroinvertebrate data and simulated streamflow for 747 subwatersheds of varying size, land use characteristics, and location;
- 4) A series of webinars, two workshops, and a technical advisory workgroup, which contributed to a stakeholder engagement process; and
- 5) A conceptual plan for a basin-wide comprehensive water resource plan, which also includes a review of existing computer-based, decision-making tools.

Flowing waters in the study area were divided into small (wadeable) and large (non-wadeable) types because the biological data collected in these two types are not comparable and require different analytical approaches for determining flow needs. Streams and small rivers are defined as Strahler order<sup>3</sup> 4 or less and wadeable. In the Middle Potomac study area, these waters correspond to the 1a, 1b, 2, and smaller 3 size categories of the Northeast Aquatic Habitat Classification System<sup>4</sup> (NEAHCS; less than 1,000 sq. mi. in upstream drainage area). Large rivers are the major tributaries of the Potomac River mainstem in the study area, as well as the mainstem itself. They correspond to NEAHCS categories 4 and greater (equal or greater than 3,861 sq. mi. in upstream drainage area).

The large river assessment approximately followed the Ecologically Sustainable Water Management (ESWM) approach described by Richter et al. (2003) which relies largely on published literature and expert judgment to identify environmental flow needs. The paucity of biological data in the study area's large rivers necessitated an ESWM approach. This study component involved (1) a literature review of basin-wide flow ecology relationships for flow-dependent species and (2) a workshop of river professionals to develop environmental flow recommendations for the mainstem Potomac River. The methods and results of the large river environmental flow needs assessment are discussed in Chapter 3.

---

<sup>2</sup> Environmental flows are defined as the seasonally variable flows of water that sustain healthy river ecosystems and the goods and services that people derive from them.

<sup>3</sup> In the Strahler stream order classification system class 1 streams are headwater streams and each successively higher number indicates a larger stream (Strahler 1957).

<sup>4</sup> The Northeast Aquatic Habitat Classification System is a standard classification system which describes stream systems across thirteen northeastern states (Olivero and Anderson 2008).

The future water use component projects the magnitude of water withdrawals, and especially consumptive withdrawals, in the Middle Potomac study area by 2030. These projections were used to complete a spatial assessment of watersheds at greatest risk of future hydrologic alteration. An additional element of this component was to compare estimates of future water use with those generated by a previous study (Steiner et al. 2000). Future water use projections are presented in Chapter 4.

The stream and small river assessment was more quantitative than the large river assessment, relying on a large macroinvertebrate database compiled from 23 local, state, and federal monitoring programs and a hydrologic model that simulated different flow conditions at locations to match the biological data. This study component of the MPRWA focused on macroinvertebrate communities because there is no fish data collected basin-wide with comparable methods. An analysis of biology, hydrology, and watershed characteristics was done using a variation of the ELOHA approach described in Poff et al. (2010). It is one of the first attempts to perform a large scale quantitative analysis using the ELOHA approach. This study component (1) identified flow metrics that, when altered, associate with strong biological response, and; (2) developed flow alteration-ecological response (FA-E) relationships that describe how aquatic ecology responds to flow alteration in a semi-quantitative manner. The relationships can be used to aid in the future development of environmental flow recommendations for tributary streams. The methods, results, and implications of the small stream environmental flow needs assessment are discussed in Chapter 5 along with lessons learned about the ELOHA approach.

Stakeholder education and engagement was an important study component. The project team solicited advice on methods and draft result interpretation, and shared results with government agencies and other parties with significant roles in water management. The stakeholder education and engagement supported an on-going dialogue among stakeholders about the sustainable use of stream flows for all purposes and contributed to state-level water management decision making processes. Stakeholder engagement is discussed in Chapter 6.

Water resources planning in the Potomac basin historically has been done by the five individual jurisdictions: the District of Columbia, Maryland, Pennsylvania, Virginia, and West Virginia. Water use and basin development in each jurisdiction affects the other jurisdictions. The fifth study component produced a concept for a basin-wide comprehensive plan process. The time and resources required to initiate and conduct a comprehensive plan are beyond what is available within this project, so this component developed a “plan” for the basin-wide comprehensive plan. Because there is no single agency with regulatory authority for water planning across the Potomac watershed, the concept for the comprehensive plan is based on a collaborative, stakeholder driven, model. The basin-wide comprehensive plan is described in Chapter 6, along with a review of computer-based, decision-making tools that have been developed by states outside the Chesapeake Bay basin.

Chapter 2 sets the stage for the major components in the project by describing existing conditions in the watershed. Chapter 7 presents overall study conclusions, lessons learned, and next-steps. Supporting materials, including datasets, additional information on methods, stand-alone reports that address aspects of the study, and copies of stakeholder outreach products are provided in technical appendices, which are referenced at appropriate places in Chapters 2-6 and are available on disc. The overall organization of the report is depicted in Table 1.

**Table 1. Organization of the MPRWA Report.**

<b>MPRWA Main Report</b>	
<b>Chapter</b>	<b>Description of Contents</b>
1	Introduction
2	Existing Conditions
3	Component #1 – Large River Environmental Flow Needs
4	Component #2 – Future Water Use Projections
5	Component #3 – Stream Environmental Flow Needs
6	Components #4 and #5 – Stakeholder Outreach and Implementation Options (including Comprehensive Water Resources Plan)
7	Conclusions

<b>MPRWA Technical Appendices (on Disc)</b>	
<b>Appendix</b>	<b>Description of Contents</b>
A	Component #1 – Large River Environmental Flow Needs Report
B	Component #2 – Water Withdrawals and Consumptive Use in the Potomac River Basin Report
C	Component #3 – Compilation of Measured Stream Data
D	Component #3 – Overview of the Application of Ecological Limits of Hydrologic Alteration
E	Component #3 – Development and Refinement of Hydrologic Model
F	Component #3 – Flow Metric Testing
G	Component #3 – Stream Classification
H	Component #3 – Basin-Wide Hydrologic Alteration Assessment
I	Component #3 – Development of Flow Alteration – Ecological Response Relationships
J	Component #4 – Stakeholder Engagement
K	Component #5 – Implementation Options


# Chapter 2. Existing Conditions

**Chapter Summary:** This chapter summarizes existing conditions in the Potomac River basin, including physiography, climate, hydrology, land uses, water withdrawals and discharges, impoundments, and water quality. Emphasis is placed on factors important to aquatic communities in streams and wadeable rivers. Many statements in the chapter are based on data and information assembled from various sources and analyzed by ICPRB for previous studies. A more detailed description of the existing conditions can be found in the attached “Potomac Basin Large River Environmental Flow Needs” report (Appendix A).



## 2.1. Physiography

The Potomac River basin is the second largest drainage basin to the Chesapeake Bay after the Susquehanna River basin. The basin covers 14,670 mi<sup>2</sup> and encompasses the District of Columbia and portions of Maryland, Pennsylvania, Virginia, and West Virginia (Figure 1). The river mainstem flows 283 miles in a generally southeastern direction from the confluence of its uppermost North Branch and South Branch tributaries to the Chesapeake Bay. Other major tributaries include the Shenandoah, Monocacy, Cacapon, Opequon, and Occoquan Rivers. The Potomac River is tidally



**Figure 2. Potomac River basin boundaries.**

Major tributaries, bioregions, and average annual precipitation are indicated.

influenced for its last 113 miles. Average discharge at the mouth is approximately 14,300 cubic feet per second (cfs).

The river crosses five major physiographic provinces which have been subdivided into Level 4 “ecoregions” by the U.S. Environmental Protection Agency (EPA)<sup>5</sup>. Ecoregions are physiographic regions “of relative homogeneity in ecological systems ... soils, vegetation, climate, geology, and physiography” (Wood et al. 1999). Aggregation of these Level 4 ecoregions into “bioregions” has proven an effective method of classifying macroinvertebrate communities in the Potomac River basin and the larger Chesapeake Bay basin (Astin 2006, Buchanan et al. 2011). Four bioregions are found in the Middle Potomac study area (Figure 2). The mountainous Ridges<sup>6</sup> bioregion is characterized by high gradient, cool, trellised streams with many riffles and active down-cutting. The Valleys bioregion, interspersed between mountain ridges, has warmer, low gradient streams. Portions of this bioregion are underlain by karst geology and have a low density of streams. The Piedmont bioregion has low to moderate gradient streams with falls, islands, and rapids. The Coastal Plain bioregion—a little of which is located in the study area—has very low gradient streams on poorly drained, alluvial sediments and streams are often poorly incised and lack a defined channel. The Atlantic seaboard fall line separates the Piedmont and Coastal Plain bioregions.

## 2.2. Climate

Located in the north temperate zone of the United States, the river basin can still experience extreme temperatures ranging from -30°F in the western mountains to 105°F in the Piedmont and low-lying coastal plain. Seasons are well-defined although changeable in spring and summer due to rapid successions of cold and warm fronts. Climate change in the Middle Atlantic region is projected to increase the average yearly temperature. Average monthly precipitation is fairly even across the year (Figure 6 in Appendix A), and summer lows in stream flow are primarily due to watershed losses to evaporation and plant transpiration between March and September. Most plant transpiration is caused by forests, which cover about 63 percent of the basin west of the fall line. Agriculture also contributes. Dry periods characterized by low and infrequent precipitation can occur at any time during the year but they reduce flow levels the most when coupled with the natural summer water losses. High flows are driven by storm events, which can occur throughout the year; they are not dependent on snowmelt. Storms usually move on a northeast track parallel to the mountain ridges and coastal zone or on an easterly track parallel to the Great Lakes and Ohio River Valley. A sizable rain shadow (the dry area on the back side of a mountainous region) caused by the Appalachian mountain system exists over the Potomac basin’s Valleys and western Ridges bioregions (Figure 2). As discussed elsewhere in this report, climate change is projected to increase the numbers of both storm events and dry periods (droughts) in the basin.

## 2.3. Hydrology

Average daily flows have been measured uninterrupted since February 1895 at the Point of Rocks gage (01638500) on the Potomac River mainstem. The gage has one of the longest time series, or

---

<sup>5</sup> [http://www.epa.gov/wed/pages/ecoregions/reg3\\_eco.htm](http://www.epa.gov/wed/pages/ecoregions/reg3_eco.htm)

<sup>6</sup> The Ridges bioregion combines the Blue Ridge province and the hilltops and mountain ridges of the Ridge and Valley province.

hydrographic records, in the United States. Approximately 96 other flow gages with hydrographic records of varying length and continuity are also located on streams and rivers in the Middle Potomac study area (Appendix C Figure 2). Flow metrics are used to evaluate the patterns of flow in streams and rivers. Flow metrics describe the magnitude, duration, frequency, and timing of different portions of the hydrograph including floods and droughts. Most are calculated from daily mean values of measured or estimated flow rate, or volume per unit time, published by the U.S. Geological Survey (USGS).<sup>7</sup>

Normal flow conditions in streams and wadeable rivers of the Middle Potomac study area are considered those found in heavily forested watersheds. The area was originally blanketed with mature stands of oaks, cedars, chestnuts, and other climax species prior to European settlement. Over the last three centuries, normal flow conditions have been altered directly with impoundments, withdrawals, and discharges and indirectly through many landscape changes in the watershed. Approximately 60 - 70 percent of the original forests were gone by the 1890s, the result of logging, mining, clear-cutting for agriculture, and catastrophic forest fires. Young forests began to reestablish in abandoned, marginal agricultural lands in the late 1800s, restoring some of the original forest's ecological and hydrological functions. The recovering trend reversed, however, when urbanization and population growth began to increase rapidly in the basin in the 1900s. Most of this growth was concentrated in the Washington DC area, leaving much of the basin rural. Future population growth is expected to spread outward from the Washington area and into the "Great Valley" between the Blue Ridge and Appalachian mountains.

There are numerous flow metrics—for example, there are 171 in the Hydrologic Index Tool (HIT) software program developed by Henriksen et al. (2006) for the USGS. Flow metrics representing all aspects of the hydrograph should be considered when investigating flow alteration-ecology (FA-E) relationships. Aquatic organisms in lotic environments exhibit many morphological adaptations and behavioral responses which allow them to cope with or avoid specific stresses associated with different flow conditions. The adaptive capabilities of an individual taxon determine to a large extent its population abundance, distribution, and resilience. The full and normal spectrum of flow conditions, from extreme low flows to large, powerful floods, helps to maintain the overall ecological integrity of streams and rivers (Poff et al. 2010). Some flow conditions may be essential to individual taxa in that they strongly affect survival or reproductive timing and potential; others may not. The large river component of this study (Chapter 3) focuses on the environmental flow needs of migratory and resident fish, freshwater mussels, and aquatic and riparian vegetation as well as tidal fresh communities in the Potomac River estuary. The stream component (Chapter 5) focuses on the environmental flow needs of benthic macroinvertebrates, the organisms most commonly and consistently monitored across Middle Potomac streams and wadeable rivers (waters in which biological samples can be collected without boats).

## 2.4. Impoundments

The Potomac is presently one of the least dam-regulated river basins in the eastern United States. The combined storage capacity of all major impoundments in the basin upstream of Washington, DC makes up less than 7 percent of median annual flow. A total of 481 impoundments are identified in the Potomac River basin by the National Inventory of Dams (2007). Many of the

---

<sup>7</sup> <http://waterdata.usgs.gov/nwis>

smaller impoundments are run-of-river and comparable in function to structures such as beaver dams. There are 153 “significant” impoundments, meaning the normal storage capacity is greater than 10 percent of mean annual flow volume and/or the dam is actively used or was once used for hydroelectric purposes (Appendix C Figure 1). In the Middle Potomac study area, the largest impoundments with respect to median annual flow volume are listed in Table 2. Included are two large impoundments that are technically upstream of this project’s study area: Jennings Randolph Lake and Savage River Reservoir in the North Branch Potomac River drainage. They are important in modeling flows in the Middle Potomac watershed. Only four of the impoundments listed in Table 2 are operated for water supply purposes. Other purposes for the selected impoundments include hydroelectric operations, recreation, and flood management. Impoundments may have ecological impacts upstream of dams in addition to downstream impacts. These within-pool non-flow related ecological impacts are not addressed in this study.

USACE owns and operates Jennings Randolph Lake for flood risk management, water supply, and recreation. Savage River Reservoir is owned and operated by the Upper Potomac River Commission for flood risk management, and water supply. Its construction was completed in 1953 by USACE under Section 7 of the 1946 Flood Control Act. USACE provides operational direction for the Savage River Reservoir, primarily for flood control in concert with Jennings Randolph Lake. Because the water from one reservoir is alkaline and acidic from the other, USACE provides a system operation on the North Branch of the Potomac River that allows for improved water quality and adequate water supply.

**Table 2. Impoundments in the Middle Potomac River study area and North Branch Potomac River holding more than 10 percent capacity of annual flow volume.**

Not included are impoundments upstream of the ones listed in the table that also have more than 10 percent capacity. Also not included are impoundments selected for inclusion in the study solely for hydroelectric purposes. Ratio: the normal storage of each impoundment, obtained from the National Inventory of Dams and converted to cubic feet, divided by model simulated median flow converted to cubic feet per year.<sup>1</sup>

Impoundment Name	Normal Storage (ac-ft)	Median Flow (cfs)	Ratio (Normal Storage to Annual Flow)
Jennings Randolph Lake <sup>2</sup>	130,900	102.27	1.77
Rocky Gap Dam	5,381	4.58	1.62
Little Seneca Lake <sup>2</sup>	13,050	13.39	1.35
Occoquan Reservoir <sup>2</sup>	25,567	53.95	0.65
Lake Barcroft	2,500	6.45	0.54
Sleepy Creek Dam	2,460	6.58	0.52
Patterson Creek No. 4 Dam	1,989	5.79	0.47
Blairs Valley Dam	486	2.12	0.32
Savage River Reservoir	20,000	90.78	0.30
Lake Gordon <sup>2</sup>	3,633	20.68	0.24
Patterson Creek No. 41 Dam	5,480	33.91	0.22

<sup>1</sup> Values for Occoquan Reservoir are observed values provided by Fairfax Water and the Occoquan Watershed Monitoring Lab. Normal storage capacity was determined from a 2010 bathymetric survey.

<sup>2</sup> Impoundments are used for water supply and their flows can be affected by dam operations.

## 2.5. Water Withdrawals and Discharges

Freshwater withdrawals in 2005 averaged 6,185 cfs for the entire Potomac River basin and represented nearly half of the estimated surface freshwater entering the estuary from all streams and rivers in the basin. Most of withdrawals (about 97.5 percent) are from surface waters. The available information suggests 78.5 percent of withdrawals are for power generation, 18.28 percent for drinking and domestic uses, 2.12 percent for industry, 0.92 percent for mining, and 0.18 percent for agriculture. Hydrologic impacts of withdrawals in the upper basin are balanced to a large degree at some point downstream by discharges which return most of the water to the river. The exception is the free-flowing Potomac River directly above the head-of-tide, where an average 574 cfs is taken from the river to supply the Washington, DC metropolitan area (WMA) and returned to the Potomac estuary at the Blue Plains wastewater treatment plant, rather than to the non-tidal river. During dry periods, these withdrawals have a large local impact on river flow but do not substantially alter the total volume of freshwater to the estuary. The Washington Aqueduct, a division of the Baltimore District, USACE, supplies drinking water for a substantial portion of the WMA, including the District of Columbia, Arlington County, Virginia, and the City of Falls Church, Virginia.

One large impact of withdrawals and discharges not related to flow alteration is their effect on water quality. As more and more people repeatedly withdraw, use, and discharge surface waters, water quality can deteriorate to the point of overwhelming the stream's assimilative capacity. Systems operated under state and federal requirements, including the Clean Water Act, can ameliorate potential water quality issues resulting from loss of assimilative capacity.

## 2.6. Land Uses

The western side of the Middle Potomac study area is mostly forested, the middle is heavily agricultural, and the eastern side intersects the heavily urban WMA (Figure 3). Deforestation and urbanization in a watershed, such as experienced in the Potomac River basin, increases the proportion of rainfall running off the landscape instead of seeping into the ground to replenish aquifers or to be taken up by plants. Increased runoff changes the magnitude, frequency, duration, and possibly the timing of stream flows. More impervious surface area makes streams flashier and gives them greater erosive power. Flow alteration impacts are not always apparent because they can be confounded by other factors normally associated with land use. Agricultural and urban land uses are often accompanied by physical disruption of stream banks and beds, either by grazing livestock or by channelization and burial under streets. Eroded top soils and sediments accumulate in stream beds. Removal of riparian buffers leads to higher temperatures and increased runoff of nutrients, herbicides, and pesticides. The presence and impacts of these other stressors on stream communities is aptly summarized in the US EPA Stressor Identification Guidance Document (2000), on the EPA Causal Analysis/Diagnosis Decision Information System (CADDIS) website ([www.epa.gov/caddis/](http://www.epa.gov/caddis/)), and in numerous text books (e.g., Ward 1992, Wetzel 2001, Thorp and Covich 2001, Merritt et al. 2008). These impacts need to be untangled from impacts that are specifically and directly related to flow alteration.




Figure 3. Land use in the Potomac River basin.

## 2.7 Ecological Community

The information in this section is partly based on background contained in the “Potomac Basin Large River Environmental Flow Needs” report (Appendix A).

### Riverine Fishes

The freshwater portions of the Potomac River basin support approximately 102 fish species, comprising 61 native upland species, 30 introduced species, and 11 diadromous species (Jenkins and Burkhead 1994). The zoogeographic history of the basin is characterized primarily by the river’s drainage to the Atlantic Ocean. This isolation from the interior continental rivers probably has reduced the overall diversity of the system compared to other river basins of the same size (Sheldon 1988). However, fish dispersal among Atlantic slope basins occurred periodically when basins were connected during low sea-level periods (i.e. ice ages) (Jenkins and Burkhead 1994). Moreover, the headwaters of the Potomac basin have “captured” stream systems from the interior Monongahela

River basin (Thompson 1939; Schwartz 1965; Cincotta et al. 1986; Hocutt 1979), presumably increasing species richness in the Potomac River basin.

Freshwater fish communities have changed substantially over time. Localized extirpations of fish taxa have been reported in the lower Potomac River basin (Starnes 2002) and probably have occurred in upstream areas. Stocking programs have also added several fish species to the River. Jenkins and Burkhead (1994) estimated that 33 percent of the fish species in the basin are introduced, a higher proportion than for other mid-Atlantic drainages. This historical absence of some popular game fishes and the presence of multiple federal fish hatcheries near Washington, D.C. may explain the relatively high level of stocking in the basin. Starnes (2002) estimated that of the 30 non-native fishes introduced into the Potomac, 22 probably remain.

Flow regimes may influence fish life history groups in different ways. For example, striped bass and white perch are influenced by flow with respect to habitat suitability during spawning and migration, and they may use flow to cue movement and spawning behaviors. Out-migrations of American eel are closely linked to high flow events and the associated turbidity. The timing of alosid migrations as well as habitat suitability for spawning and overwintering are affected by flow. Flow regimes influence non-migratory fishes primarily through their influence on local habitat quality. For example, smallmouth bass recruitment can be limited by high flow events that scour nests, reducing recruitment.

### Macroinvertebrates

Stream macroinvertebrates are a diverse group of organisms with many morphological, behavioral, and feeding adaptations for life in flowing waters. They consume detritus, algae, bacteria, and microscopic animals, and are food for fish and birds. They have a broad range of tolerances and sensitivities to different stressors and anthropogenic pollutants, and fill many ecological niches. This makes them ideal as indicators of the health of most streams and rivers.

Aquatic macroinvertebrates are a diverse group representing numerous phylogenetic taxa, but most communities in free-flowing waters are dominated by insects. Insect life cycles are relatively short—on the order of weeks to a few years—and their responses to extreme but infrequent events (extreme droughts, extreme floods) occur in the year of the event and then fade as successive generations recover. An appropriate suite of flow metrics for macroinvertebrates may be those that reflect flow alteration occurring on a regular basis. Macroinvertebrates also are sensitive to environmental stressors that do not alter flow *per se* but that can be associated with flow and can confound the effects of flow alteration, such as water chemistry (pH, conductivity, dissolved oxygen, nutrients) and habitat condition (loss of riparian zone vegetation, sedimentation, channelization). Macroinvertebrate communities in the Potomac Basin are assessed using a uniform scoring approach with a suite of biometrics. Macroinvertebrate communities have been assessed from very poor condition to excellent, depending on the environmental stressors. The poorest stream indexes occur in highly urbanized watersheds such as those in the Washington D.C. metropolitan area. Lower scores are also present in areas with intense agricultural activity. The highest scores are typically found in minimally disturbed watersheds with low levels of pollution and stable in-stream and streamside habitats. These watersheds tend to be clustered in forested areas along the western side of the Chesapeake Bay basin (Buchanan et al. 2011).

### **Riverine Mussels**

There are 14-15 native mussel species, depending on taxonomic classification, which are recognized in the Potomac River basin, and encompass a variety of traits related to habitat use and reproduction. As with fishes, the Potomac River basin generally supports fewer freshwater mussel species than rivers draining into the interior basins of the United States (e.g. Mississippi River) (Ortmann 1913; Taylor 1985). In Atlantic-slope basins, mussel communities suggest colonization among rivers, perhaps due to fish host dispersal (Sepkoski and Rex 1974).

Potomac basin mussels exhibit considerable variation in their tolerance of flow velocities. Several species do not exhibit flow preference but are found in slack water, slow, moderate, and fast velocities. Conversely, five species occur within a single flow-velocity category, primarily slack-water or slow-water specialists. No species were fast-water specialists, but seven species occur in fast-water as well as slow-water.

Flows also affect mussels indirectly through influencing behavior of fish hosts for glochidia dispersal. Experimental trials have revealed fish-host relations for several mussel species found in the Potomac Basin. In some cases, amphibians may also serve as hosts for glochidial dispersal (Watters and O'dee 1998). Fish movements are often linked to variation in flow regimes (Fausch et al. 2001) and fishes of the mid-Atlantic highlands exhibit distinct dispersal-grain signatures (Hitt and Angermeier 2008).

### **Riparian Vegetation**

Floodplain forests and other riparian vegetative communities have varying flow regime requirements which, in un-regulated rivers, are principally factors of elevation. Ecological gradients develop from the fully inundated river bottom to frequently flooded river benches to occasionally flood-affected hill top. Within those gradients, temporal variation is important to maintain ecological complexity, where floods and droughts have differential effects, each adversely affect some communities while benefitting others. For instance, submerged aquatic plants in the Potomac experience their greatest growth and reproductive potential during years with lower flows, especially if flows are low during the growing season due to increased water clarity, as less sediment and nutrients are running off the landscape and there is greater substrate stability. Conversely they tend to grow poorly during wet summers when flows are more turbid and erosive. Up from the river's edge, however, flooding-caused tree falls promote diversity by providing openings in the canopy and opportunities for pioneer and understory species that do not occur during dry years. Floodplain plants depend on floods for seed dispersal, deposition of sediment to maintain floodplain surfaces and enrich soils, removal of debris and potential competitors from germination sites, and to provide adequate moisture conditions for germination and growth (Dixon 2003). Duration and frequency of floods upon different fluvial landforms was found to be the most important factor determining riparian vegetation communities along a river in Virginia (Hupp and Osterkamp 1985). Similarly, species richness of riparian plants increases with topographic complexity of the floodplain (Everson and Boucher 1998).

# Chapter 3. Large River Environmental Flow Needs

---

**Chapter Summary:** The large river environmental flow needs assessment is the first component of the MPRWA. The complete report, including the workshop participants' comments, can be found in Appendix A. This chapter summarizes the assessment's key findings and recommendations. The assessment was made using the Ecologically Sustainable Water Management (ESWM) approach, which relies largely on published literature and expert judgment to identify environmental flow needs. This study component produced (1) a literature review of basin-wide flow ecology relationships for flow-dependent species and (2) environmental flow recommendations for the mainstem Potomac River. The taxonomic groups investigated were fish, mussels, instream and riparian plants in the free-flowing large rivers and fish, macrobenthic invertebrates, zooplankton, and phytoplankton in the tidal fresh estuary. There was particular emphasis, driven by stakeholder interest, in low flows. The project team found no discernible adverse impact on key biotic taxa due to human modification of *flow* in the free-flowing large rivers. The team recommended that the current large river flow regimes be maintained, particularly at low flows. They concurred with the 100-mgd (million gallons per day) and 300-mgd minimum flow requirements for Little Falls and Great Falls, respectively. A diverse group of experts reviewed the study's findings and conclusions at a two-day workshop in September 2010.



## 3.1. Introduction

The assessment of large river environmental flow needs for four mainstem Potomac River segments and two large tributaries was a stand-alone component of the MPRWA. There is insufficient comparable biological data for large river segments in the Potomac to develop quantitative flow-ecology relationships using the methodology employed for smaller streams. For this reason the hydrologic needs of flow-dependent species and communities in four mainstem Potomac segments and two selected large tributaries deemed to be at risk of hydrologic alteration were identified using a modification of the Ecologically Sustainable Water Management (ESWM) approach described in Richter et al. (2003). The methods and results of that assessment are summarized here and reported in full in a stand-alone report, "Potomac Basin Large River Environmental Flow Needs" (Cummins et al. 2011), which is included in Appendix A. The National Park Service (NPS) provided funding for this portion of the MPRWA.

The Potomac Basin Large River Environmental Flow Needs assessment was developed by a research team from the ICPRB, TNC, Leetown Science Center Aquatic Ecology Branch of USGS, and the Potomac Environmental Research and Education Center of George Mason University (GMU). It includes a comprehensive literature review, development of flow hypotheses, assessment of large river environmental flow needs, statistics proposed to track those flow needs, and recommendations for additional research, monitoring, and analysis to improve understanding of flow needs. As part of the literature review, more than 480 sources of information were collected, reviewed, and organized into a searchable on-line database (Appendix G in Cummins et al. (2011)).

The Potomac Basin Large River Environmental Flow Needs assessment included a workshop, held September 22-23, 2010, at the National Conservation Training Center in Shepherdstown, WV (hereinafter referred to as the September 2010 workshop), at which 60 hydrologists, biologists, engineers, water resource managers, and regional and national experts on flow and river ecology discussed draft findings from the project research team. A summary of that workshop and participants' comments is available as Appendix H in Cummins et al. (2011). At the September 2010 workshop, participants concluded that despite the detailed review and analysis of currently available literature, more research and monitoring is needed to fill information gaps to define ecologically protective flow alteration thresholds for Potomac large river systems. A new technical working group will be convened to plan for additional hydroecological research and monitoring that will support the development of quantitative large river flow recommendations.

### **3.2. Large River Flow Needs Assessment Approach**

A preliminary risk assessment of sub-basins in the Potomac River Basin was conducted to identify watersheds with the highest risk of hydrologic alteration from multiple factors. To this end, 35 sub-basins and 5 Potomac River mainstem segments were assessed. The risk assessment methodology consisted of four phases: (1) identifying possible risk factors in the Potomac Basin related to hydrologic alteration, (2) identifying correlation of risk factors with other risk factors and with Indicators of Hydrologic Alteration (IHA) metrics (Richter et al. 1996), (3) establishing risk thresholds utilizing hydrologic metrics, and (4) calculating a cumulative risk index for each sub-basin and mainstem segment based on selected risk factors.

Based on the results of this hydrologic alteration risk assessment, four Potomac mainstem segments and two large tributary streams were selected for this flow needs assessment due to the high count and severity of risk factors that can lead to altered hydrology (Appendix B in Cummins et al. (2011)). These river segments (Figure 1) were:

- 1) Potomac mainstem from the confluence of the Shenandoah River to Point of Rocks;
- 2) Potomac mainstem from Point of Rocks to Great Falls;
- 3) Potomac mainstem from Great Falls to Chain Bridge (Potomac Gorge or Fall Zone);
- 4) the tidal fresh Potomac estuary from Chain Bridge to Occoquan Bay;
- 5) Monocacy River mainstem; and
- 6) Opequon Creek mainstem.

The Potomac Gorge is of special concern because of its relatively unique and rare riparian biological communities. One charge to the study's research team was to re-examine the 100 million gallons per day (mgd; 155 cfs) minimum flow-by requirement for the Potomac Gorge. This flow-by value was established by the Maryland Department of Natural Resources' (MDDNR) Potomac River Environmental Flow-by Study in 1981, as a follow-up to the 1978 Potomac River Low Flow Allocation Agreement which identifies the DC area water supply allocations during low flow periods.

The research team did not find sufficient research or empirical data to define thresholds of ecologically acceptable hydrologic change for the large rivers of the Middle Potomac River study area. Therefore, the ESWM approach was applied. Four plant communities, twelve fish species, and sixteen native mussel species were identified and used to represent the diversity of species, the flow-

ecology relationships, and the flow needs of communities found in the large, free flowing rivers of the basin. The research team used the available literature and professional judgment to develop four general flow-ecology hypotheses that apply to a broad range of species/communities and 18 specific flow-ecology hypotheses tailored to selected indicator taxa in the non-tidal Potomac large river segments and selected large tributaries (see below and Cummins et al. (2011), pp. 50-51).

Phytoplankton, aquatic grasses, zooplankton, and benthic invertebrate communities, and four fish species were used to represent key aspects of tidal freshwater ecology and its responses to low freshwater flows in the tidal fresh estuary. In general, freshwater inflow to the estuary delivers nutrients, sediment, and other constituents, and influences the salinity gradient which governs structure and function of biological communities along the entire length of the estuary. Low flow effects on estuarine biota are for the most part indirect and realized as a change in salinity, or the volume proportions of fresh and salt water. Flow alteration as a factor affecting the Potomac tidal fresh biological communities is presently far outweighed by the effects of poor water quality and other stressors. Although of lesser importance than water quality impacts, seven general flow-ecology hypotheses for the tidal fresh estuary also were articulated (see below and Cummins et al. (2011), p. 71).

---

## Flow-Ecology Hypotheses for Potomac Nontidal River Communities

### General hypotheses

- 1) Species richness will peak at intermediate levels of flow variability (*sensu* Connell 1978).
  - a. Too many low-flows will extirpate riverine biota due to a cascade of flow-induced effects on water quality, connectivity, biotic interactions (i.e., predation and competition).
  - b. Too many high-flows will extirpate some riverine biota through sheer-stress effects and habitat loss.
  - c. An “intermediate” level of flow variability will increase riverine species richness by creating habitat features and limiting competition.
- 2) Low and high flow effects will be mediated by the spatial proximity and abundance of flow refugia and organismal vagility.
- 3) The mechanisms of flow-effects will vary across spatial and temporal scales.
  - a. Spatial
    - i. At local-scales (i.e., within 1-10 mile-long river reaches), flows affect riverine biota primarily through physiological and behavioral pathways.
    - ii. At regional-scales (i.e., more than 10 mile-long river reaches), flows affect riverine biota primarily through recruitment and metapopulation dynamics.
  - b. Temporal
    - i. Over short time periods (i.e., hours-days), flows affect riverine biota primarily through physiological and behavioral pathways (e.g., sheer stress).
    - ii. Over long time periods (i.e., years-decades), flows affect riverine biota primarily through habitat-forming processes (e.g., substrate organization and mesohabitat structure).
- 4) Behavioral, phenotypic, and physiological species traits will predict organismal and population sensitivity to flow regimes.

## ***Specific hypotheses***

### Plant Communities

- 5) Submerged aquatic plants experience their greatest growth and reproduction during years with lower flows during the growing season due to increased water clarity and greater substrate stability.
- 6) Floodplain plants depend on floods for seed dispersal, deposition of sediment to maintain floodplain surfaces and enrich soils, removal of debris and potential competitors from germination sites, and to provide adequate moisture conditions for germination and growth.
- 7) Flooding-caused tree falls promote diversity by providing openings in the canopy and opportunities for pioneer and understory species that do not occur during dry years.
- 8) Duration and frequency of floods upon different fluvial landforms is the most important factor determining riparian vegetation communities.
- 9) Species richness of riparian plants increases with topographic complexity of the floodplain.

### Fishes

- 10) Out-migrations of American eel (*Anguilla rostrata*) and alosids are triggered by high-flows and associated water quality conditions (i.e., turbidity).
- 11) Fishes exhibiting K-selected reproductive strategies with fewer, longer-living offspring will be more vulnerable to stochastic high-flows and floods than R-selected species with many shorter living offspring (e.g., margined madtom, *Noturus insignis* versus white sucker, *Catostomus commersoni*, respectively).
- 12) Pelagic fishes will be influenced by stochastic flow variability more than benthic fishes.
- 13) Fishes exhibiting simple lithophilic spawning strategies (i.e., no parental care, gravel-spawning species) will be influenced by stochastic flow variability more than other reproductive strategies.
- 14) Riverine fishes exhibit a bimodal distribution of body sizes in response to natural flow regimes (i.e., regulated rivers will exhibit unimodal distributions).
- 15) Spring peak-flows regulate smallmouth bass (*Micropterus dolomieu*) recruitment.
- 16) Bedrock-dominated river reaches will be more prone to high-flow extirpations than freestone dominated river reaches (i.e., microhabitat refugia).
- 17) River reaches containing stream-river confluences will be less-subject to flow-induced extirpations than river reaches without such stream network connectivity (i.e., macrohabitat refugia).

### Mussels

- 18) Winter flow conditions will influence recruitment in long-term brooding mussel species (represented by *Lampsilis* sp.) more than in short-term brooding species (represented by *Elliptio* sp.).
- 19) Recruitment in short-term brooding mussel species will be influenced by stochastic effects of peak-flows more than long-term brooders.
- 20) Deep-water mussel species will be less subject to drought than shallow-water species.
- 21) Mussel populations exhibit patch dynamics at the meso-habitat scale such that isolated riffles are more vulnerable to flow-induced adverse effects than “connected” riffles.
- 22) Mussel fish host generalists are less subject to flow-induced adverse effects than fish host specialists.

## Flow-Ecology Hypotheses for Biological Communities in the Tidal Freshwater Potomac Estuary

- 1) A range of freshwater flows gives intermittent opportunity to both freshwater and brackish water species and produces a biologically diverse estuarine ecosystem.
  - 2) Salinity, and specifically the location of the “salt wedge,” is a surrogate measure of freshwater flow in the upper, tidal fresh estuary.
  - 3) The flow “needs” of most freshwater species in the tidal fresh segment are typically a reflection of their salinity preferences and tolerances.
  - 4) High flows, in conjunction with the daily light cycle, temperature and/or turbidity, cue diadromous fish migrations into estuaries and rivers in spring and out-migrations in autumn.
  - 5) Eutrophication and sedimentation of the Potomac River have changed many estuarine flow-ecology relationships:
    - a. The tidal fresh reach of the estuary is irreversibly longer and shallower than it once was;
    - b. Wastewater returns to the tidal fresh estuary result in increasing rather than decreasing nutrient concentrations when flows from the upper basin are low; high flows from the upper basin dilute tidal fresh nutrient concentrations;
    - c. Water clarity is poorer during low flow periods rather than high flow periods;
    - d. Phytoplankton and zooplankton blooms now occur predominantly in summer rather than spring and are not as closely linked to high flows in spring as they once were;
    - e. Uneaten phytoplankton sink and cause bottom layer hypoxia and anoxia in brackish waters that can both stress and block migratory life stages of bottom-oriented fish and benthic invertebrates to the tidal freshwater;
  - 6) Depending on water quality conditions, a persistent loss of freshwater flow (e.g., due to consumptive use or increased forest evapotranspiration) could reduce the extent of tidal fresh communities relative to brackish water communities.
  - 7) Depending on water quality conditions, a persistent increase of freshwater flow (e.g., due to climate change) could increase the extent of tidal fresh communities at the expense of brackish water communities such as oyster reefs.
- 

### 3.3. Findings of the Large River Flow Needs Assessment

Drawing upon the literature review of aquatic species’ flow needs and analysis of the historical record of flows for these rivers, the following key considerations were synthesized. These considerations shaped the project team’s findings regarding large river flow needs:

- 1) The Potomac River has only minimal flow regulation and that occurs only at very low flows. There are no dams regulating flow on Opequon Creek or Monocacy River. Magnitude, frequency, and duration of high and mid-range flow events, therefore, are more influenced by land use management than operational management.
- 2) The observed river flow characteristics appear to be primarily the result of weather, climate, and land use factors, except for low flows from Great Falls to Little Falls, and potentially in the Monocacy.

## Middle Potomac River Watershed Assessment

- 3) Evidence suggests that there have been changes in flow distributions over the past 100 years, but additional analyses are required to determine the roles of climate, land use, or other factors, in those changes.
- 4) Intra- and inter-annual variability in flows is high for these stream reaches.
- 5) For aquatic species, very few studies in the literature provided directly applicable quantitative measures of flow needs (beyond velocity requirements at the individual organism scale). These requirements could not be translated to stream discharge values. The literature and expert judgment did provide qualitative descriptions of flow needs.
- 6) No research or monitoring evidence of species impairment due to current levels of flow management was found in Potomac large rivers.
- 7) Low flows in the Great Falls to Little Falls reach are lower than they would otherwise be due to drinking water withdrawals at, and above, Great Falls. Minimum flow-by's of 300 mgd (464 cfs) at Great Falls and 100 mgd (155 cfs) at Little Falls were recommended by MDDNR (1981) which have been maintained by the Washington metro region water suppliers under the guidance of the ICPRB CO-OP section since the early 1980s. During that time flows have rarely approached these limits. In 2002, when flows were approaching these levels, field observations in surveyed areas did not identify any stressed communities, and there did not seem to be a significant loss of habitat in these reaches.
- 8) The flow “needs” of most freshwater species in the tidal fresh river segment are typically a reflection of their salinity preferences and tolerances. High river flows can benefit taxa and life stages that prefer freshwater while low flows can benefit taxa and life stages that prefer salt water.
- 9) Eutrophication and sedimentation of the tidal Potomac River have significantly changed many estuarine flow-ecology relationships. The flow needs identified for tidal freshwater biota do not consider the very significant confounding influence of the tidal freshwater Potomac River's poor water quality. Nor do they consider the flow needs of higher salinity taxa such as oysters, young-of-year menhaden, and older, resident striped bass.
- 10) Future impacts on flow from climate change are uncertain, but studies have suggested that impacts in the Middle Atlantic region of the U.S. will be lower in magnitude than elsewhere and may result in both greater precipitation and higher temperatures (which could increase demand for electricity and consumptive water use).

Considering these points, the team's approach was less a question of determining what flows are required to *restore* these river sections, and more a matter of *defining and characterizing how existing flows are functioning* to maintain ecological values. The Large River Flow Needs assessment provides that characterization in both qualitative and quantitative terms.

### 3.4. Conclusions for Large River Flow Needs Assessment

Based on the considerations described in Cummins et al. (2011) and summarized above, the project team made the following conclusions to characterize the Potomac's large river flow needs:

- 1) For the entire range of flows, the current flow characteristics, as defined in Cummins et al. (2011)<sup>8</sup>, should be maintained as a precautionary principle. If additional monitoring and analysis provides more definitive indications of biological degradation due to flow, then other more protective flow recommendations might be needed.
- 2) Extreme floods: High flows and floods in the river segments in this study are not controlled by dams or other structural measures and so there are no operational mechanisms for controlling high flows. The impact on extreme high flows of impervious surface area and extent of vegetative cover in the watershed upstream of these river segments is not known presently but is being evaluated as part of the Middle Potomac River Watershed Assessment.
- 3) Small Floods: No observed major problems, so current flow characteristics should be maintained.
- 4) Low Flows at Potomac Harpers Ferry to Point of Rocks: This section benefits from slightly augmented flows during low flows due to water quality and water supply releases from Jennings Randolph and Savage River reservoirs. There are no observed flow-related, ecological problems in this reach, therefore, recommend maintaining current flow characteristics.
- 5) Low Flows at Potomac Point of Rocks to Great Falls: Withdrawals should be managed so that Potomac River flows do not fall below those experienced in the 1999 and 2002 droughts. It is recommended also that a stream flow gage be installed to measure actual flow levels at the Great Falls weir.
- 6) Low Flows at Potomac Great Falls to Little Falls: (a) prior (1981) recommendation for a 300 mgd minimum flow should be continued, but (b) implement an ecological monitoring program to better understand if there are impacts and need to adapt our management, and (c) as a precautionary measure until this study is completed, develop reservoir operating procedures which give consideration to maintaining variability at extreme low flows.
- 7) Low Flows at Potomac Little Falls to Chain Bridge (tidal river): (a) maintain the existing 100 mgd minimum flow-by, but (b) implement an ecological monitoring program to better understand if there are impacts and need to adapt our management, and (c) as a precautionary measure until this study is completed, develop reservoir operating procedures which give consideration to maintaining variability at extreme low flows.
- 8) Low Flows at Potomac Chain Bridge to Occoquan Bay: Water quality is the major determinant of biological health, not freshwater flow. Current flow characteristics should be maintained.
- 9) Low Flows at Monocacy River and Opequon Creek: As a conservative measure, until additional investigations of potential low flow impairment can be conducted, current low flow statistics should be maintained and withdrawal volumes not be allowed to push flows below those observed in 1999 and 2002.

---

<sup>8</sup> Table 16 in Cummins et al. (2011) provides numerical values for 20 flow statistics for stream gages on Opequon Creek, the Monocacy River, and at Point of Rocks and Little Falls on the Potomac. For the Little Falls gage, statistics are provided for adjusted (water supply withdrawals added to observed flow) and unadjusted flows.

### 3.5. Large River Information Gaps and Monitoring Needs

For most of the species discussed in the Potomac Basin Large River Environmental Flow Needs assessment, existing information is adequate only for qualitative estimates of how normal variability in population and distribution is affected by alterations in flow conditions. Therefore, additional monitoring, or analysis could help to define ecologically acceptable levels of hydrologic change, or acceptable thresholds of hydrologic alteration from current conditions for species in large rivers.

Although the September 2010 workshop aimed to define the full range of natural flow conditions for six river segments, the participants paid greatest attention to low flow conditions and ecological response in the Great Falls to Little Falls river segment. Some of the workshop participants' most significant suggestions include:

- 1) Address monitoring and data analysis gaps identified in the 2004 and 2005 Potomac low flow workshops, including studies to better understand "normal" variation of species populations and ranges, and studies to better understand the effects of extreme low flows on species and their habitat.
- 2) Monitor effects of high flows on floodplain plants and communities. The Potomac Gorge should be a priority site for monitoring the effects of high flows due to its great concentration of rare flood-dependent vegetation community types.
- 3) Monitor impacts of low flows on mussels. The species recommended are *Elliptio complanata*, *Pyganodon cataracta*, *Utterbackia imbecillis*, *Lampsilis sp.*, and possibly *Strophitus undulatus* and *Alasmidonta undulata*.
- 4) Monitor fish to establish a better quantification of their flow needs, including fall young-of-year fish, alosid passage over the Little Falls weir, and in- and out-migration of fish. Also pursue research on fish species which live near drinking water intake pipes, focusing on short rather than long life span species.
- 5) Acknowledge opportunities and limitations for researching flow-ecology relationships in other species groups:
  - a) Macroinvertebrates may be useful for flow-ecology research, but large river study protocols are not well developed. Crayfish may be an important group to study as they are an important food source to other species, and to discern their life-cycle relationships to flow.
  - b) Amphibians and reptiles are difficult to study because they are mobile, but could be of interest for tracking loss of habitat if flooding is reduced.
  - c) Cormorants are important as fish predators, but they are mobile, part-time residents, and population changes may be due to factors other than river flow and fish (prey) abundance.
- 6) Track cumulative upstream consumptive use of water because of its potential role in reducing extreme low flows.
- 7) Investigate the use of remote-sensed imagery, such as Light Detection and Ranging (LIDAR) high-resolution topographic data, for determining the extent of loss of habitat at different flow levels.
- 8) Consider pursuing a modified Instream Flow Incremental Methodology (IFIM) study for evaluating the relationship between flow and habitat at flows below 1000 mgd in the stretch from Great Falls to Little Falls.

## Middle Potomac River Watershed Assessment

September 2010 workshop participants recognized that funding, staff time, and public attention or political will are constraining factors for developing a large river flow needs research and monitoring program. They concluded that a coordinated federal, interstate, and academic partnership would be needed to obtain resources and long-term commitment to: (a) developing a baseline during mid-range flow conditions, and (b) enabling monitoring and additional research during the more extreme high and low flow conditions.

# Chapter 4. Future Water Use Projections

---

**Chapter Summary:** The second component of the MPRWA project examined the expected magnitude of future increases in water withdrawals and consumptive losses, to identify the principal causes of those increases, and to estimate how those increases might affect the stream flow regime. These future projections can be used to determine how future flow alterations might influence biological communities as an element of the stream flow needs assessment, and to complete a spatial assessment of watersheds at greatest risk of future hydrologic alteration. The future water use component of this study involved a comparison of estimates of future water use with those generated by a 2000 study of basin-wide (above Little Falls) consumptive use (Steiner et al. 2000). The methods and assumptions used to estimate increased water use, by use sector, and the comparison to the 2000 basin-wide consumptive use study are fully described in a report titled “Water Withdrawals and Consumptive Use in the Potomac River Basin.” The report is a stand-alone component of the MPRWA; it is included in this report as Appendix B. The application of future water use scenarios to estimate impacts on the stream flow regime, as represented by alteration in selected flow statistics, is described in Appendix E.



## 4.1. Methods for Future Water Use Projections

Current water withdrawal (surface and groundwater) estimates were based on USGS county-level withdrawal data attributed to specific withdrawal points obtained from each basin state with 2005 as the base year. Withdrawals were assigned to one of six use sector categories: (1) domestic and public supply, (2) mining, (3) thermo-electric power, (4) industry, (5) livestock, and (6) irrigation. Withdrawals in each water use sector were assigned a consumptive use factor based on historical values assigned by the USGS. The methods and data sources used to project future water use varied by use sector. Projections were made to 2030. The assumptions used to predict future consumption become less reliable as the projection time period increases. Therefore, the analysis is constrained to forecasted conditions for 2030.

Six scenarios were designed to provide insight into possible flow impacts across a range of possible future conditions (Appendix B, section 5). Rates of change in withdrawals for the mining, thermo-electric power, industry, livestock, and irrigation sectors were constant across the scenarios unless otherwise noted. The scenarios are:

- 1) DP1 – 0 percent per year increase in per capita withdrawal in domestic and public supply, so the increase in total withdrawals is due solely to population growth.
- 2) DP2 – 1.82 percent per year increase in per capita withdrawal in domestic and public supply, coupled with population growth.
- 3) DP3 – 4.38 percent per year increase in per capita withdrawal in domestic and public supply (referred to as “base” in Appendix B), coupled with population growth.
- 4) Power – New closed cycle, thermo-electric power plant on the Monocacy River in Frederick County, MD, and conversion from open cycle to closed cycle cooling at the other thermo-

- electric power plants. Closed cycle cooling increases consumptive use but dramatically reduced withdrawals. This scenario builds on the DP2 growth rate for domestic and public supply.
- 5) Climate Change – Applies Intergovernmental Panel on Climate Change (IPCC) projections for temperature increase and consequent changes in irrigation, power demand, and summer demand for domestic and public supply. This scenario builds on the DP2 growth rate for domestic and public supply.
  - 6) Hot and Dry – Domestic and public supply sector withdrawals increased by 15.2 percent during April through August. Power sector withdrawals increased by 6.15 percent during May through September. Irrigation sector withdrawals increased by 284 percent during May through September. This scenario builds on the DP2 growth rate for domestic and public supply.

The six scenarios were developed as representative of future conditions of interest for basin stakeholders based on known development trends, projected population trends, or projected weather and climactic conditions. The DP3 scenario is characterized as the “base” scenario on which the other five scenarios are derived. The DP2 scenario corrects data inconsistencies in several of the withdrawal values reported by USGS for certain jurisdictions. The resulting changes led to a reduction in the annual change in the per person withdrawal rate. The DP1 scenario isolates the growth in withdrawals in the domestic and public supply sector that was due solely to projected population growth. This scenario assumed that there was no change in withdrawal per person over time, holding the withdrawal per person constant at the 2005 rate. The power sector scenario assesses the withdrawal and consumptive use impacts that could be expected if a new power plant becomes operational in the basin as well as assessing potential impacts of retrofits at existing power plants. This scenario is based on current expectations of power plant development in the basin. The climate change scenario assesses the impact of climate changes on the 2030 water demand projections in the DP2 scenario. The hot and dry scenario examines water withdrawals under weather conditions that are occasionally historically experienced in the basin, not factoring in changes to incidence of hot and dry weather that might be altered under future climate change conditions.

An overall increase in water withdrawals is forecasted for all future scenarios. The scenarios assume there will be no decrease in future water demand in the domestic and public supply sector. This is predicated on the assumption that past usage indicates future usage (per capita), and that conservation measures mandated by the Energy Policy Act of 1992 have largely been implemented and energy saving reductions realized. Changes in plant operations at thermo-electric power plants (conversion to close-cycle) are assumed to decrease withdrawals but increase consumptive use. The future scenarios were designed conservatively, to ensure that planning efforts do not result in overutilization of water resources. A detailed description of the methods and calculations used to project future water uses can be found in Appendix B.

These scenarios, excluding the DP3 scenario, were used to configure future hydrologic model scenarios with which flow time series were generated, flow statistics calculated, and flow alteration determined. Future flow time series were generated for 153 subwatersheds representing the Chesapeake Bay Program’s Hydrologic Simulation Program-Fortran (HSPF) model segments in the study area. Future flow alteration is the relative difference between baseline and future scenario flow metrics. The methods for modeling the future scenarios and maps of predicted future hydrologic alteration for selected flow metrics are presented in Chapter 5 (Figures 20a-d) and in Appendix E (FutureScenarios\_011912.pdf).

## 4.2. Findings for Future Water Use Projections

Table 3 shows cumulative withdrawals and consumptive use, by use sector, for the Middle Potomac River study area and North Branch watershed combined<sup>9</sup> (Figure 1). The North Branch is included in estimates for the MPRWA study area because it is upstream of the study area. See Appendix B for details on source information and methods for forecasting changes in each use sector. There are uncertainties in the estimates for each use sector.

The use sector with the largest water withdrawals was power generation, accounting for 68 percent of total withdrawals in the base year 2005 and 42 - 65 percent of total withdrawals in the six future scenarios. Domestic and public supply was the next largest sector, accounting for 27 percent of total water withdrawals in 2005 and 29 - 55 percent in the future scenarios. The combined agriculture, industry, and mining sectors accounted for 5 percent of total withdrawals in 2005 and 3 - 5 percent in the future scenarios.

Results for consumptive use are significantly different from withdrawals. The domestic and public supply sector accounts for 84 percent of total consumption in 2005 and 84 - 94 percent in the future scenarios. The power sector is a distant second (8 - 12 percent), and agriculture, industry, and mining each account for 2 percent or less of total consumption. One reason for the very high consumption rate in the domestic and public supply sector is the way consumption is calculated. If withdrawals eventually are returned to free-flowing waters, a consumption rate of 11 percent is applied. If, however, withdrawals from free-flowing waters eventually are discharged to an estuary, as they are in the parts of the Washington D.C. metropolitan area, they are considered 100 percent consumed. In the estuary, the discharged water is no longer considered available for domestic and public supply withdrawals and is therefore a computational “loss” included in the consumptive use total. In the study, all of the water withdrawn from the Potomac River mainstem by Washington Aqueduct, Washington Suburban Sanitary Commission, and the City of Rockville is assumed to be discharged to the estuary, primarily through the Blue Plains Wastewater Treatment Plant. Except for the reaches between withdrawal and discharge locations, discharged water is not truly lost from the river system; it represents a very large freshwater input to the estuary.

At the scale of the MPRWA study area, different assumptions or calculation methods for use sectors other than domestic and public supply are not likely to have much impact on estimate of total consumptive use. Agriculture does become a more significant water consumer during periods of drought. The hot and dry scenario, roughly equivalent to the 1930-31 drought of record, increased agricultural consumptive use from 5 mgd in 2005 to 13 mgd in 2030. Agriculture sector water use in other scenarios decreases from 2005 to 2030 due to a predicted decrease in the agricultural acreage (also present in the hot and dry scenario). Consumptive use is higher in the hot and dry scenario and climate change scenario compared to the DP2 scenario from which they were built. In the power sector, there is an overall increase in consumptive use in all future scenarios due to population growth. In the power sector of the power scenario, withdrawals and consumption reflect the change to closed cycle technology, which withdraws less water but has a much higher consumptive rate per kilowatt hour of electricity produced.

---

<sup>9</sup> Tables with equivalent information for the entire Potomac basin (Potomac) and for the upper basin above the fall-line (AFL) are available in Appendix B supplemental tables. Also included are county breakouts for each scenario.

**Table 3. Withdrawals (mgd) and consumptive use (mgd) in the Middle Potomac River Watershed Assessment study area (includes the North Branch Potomac River watershed), by sector, for six future scenarios.**

See text for full descriptions of scenarios. The values presented are rounded to the nearest whole number. Comparable tables for the entire Potomac River basin (Potomac) and the upper basin above the fall-line (AFL) are provided as supplemental tables in Appendix B. In this table, the livestock and irrigation use sectors are combined into agriculture.

Scenario		Agriculture	Domestic & Public	Industry	Mining	Power	Total	
<b>2005</b>								
Current	Withdrawals	7	609	84	28	1,524	2,252	
	Consumption	5	331	8	4	46	394	
								%Change from 2005
<b>2030</b>								
DP1	Withdrawals	5	856	119	30	1,895	2,904	29%
	Consumption	4	411	11	4	57	488	24%
DP2	Withdrawals	5	1,343	119	30	1,895	3,391	51%
	Consumption	4	645	11	4	57	722	83%
DP3	Withdrawals	5	2,494	119	30	1,895	4,543	102%
	Consumption	4	1,199	11	4	57	1,275	224%
Hot and Dry	Withdrawals	16	1,433	119	30	1,947	3,545	57%
	Consumption	13	688	11	4	58	775	97%
Climate Change	Withdrawals	8	2,563	119	30	1,946	4,665	107%
	Consumption	6	1,231	11	4	58	1,311	233%
Power	Withdrawals	5	1,343	119	30	1,657	3,154	40%
	Consumption	4	645	11	4	138	803	104%

Maps of future hydrologic alteration predicted in five of the six scenarios for selected flow metrics in 153 large subwatersheds are presented in Appendix D (Figures 15a-d). The direction of future hydrologic alteration in a flow metric often differed between watersheds due to differences in anticipated watershed changes. (The relationships of watershed drivers to alterations in the suite of selected flow metrics are described in detail in Appendix H.) Overall, the cumulative impacts of predicted land and water use changes coupled with the meteorological stressors in the climate change and hot and dry scenarios result in the most hydrologic alteration. The DP1, DP2, and power scenarios result in more moderate alteration because temperature, precipitation, and evapotranspiration regimes are assumed to remain constant. While the current (2005) and projected consumptive use in sectors other than domestic and public supply have a very small impact on the total when summed across the Middle Potomac watershed, consumptive use in subwatersheds may have substantial impacts. These subwatershed impacts are discussed in more detail in Chapter 5 and Appendix D. Water conservation and demand reduction should be pursued as part of a wise use of water. The scenarios described here are not prescriptions for future action.

### 4.3. Comparison to Previous Studies

A previous study conducted by the ICPRB Section for Cooperative Water Supply Operations on the Potomac (CO-OP; Steiner et al. 2000) estimated current and future demands, consumptive use rates, and available resources through 2030. Comparison of the forecasts of withdrawals and consumptive use by the previous study and by this project are difficult because of differences in data and methods; therefore, several modifications were made to the assumptions in this study to perform a direct comparison of the results.

One approach used in Steiner et al. (2000) was to provide consumptive use forecasts by HUC 8 watershed. The study area was the area above Little Falls and it did not include the WMA water utilities. The results represented use in a typical stream flow year and the approach was able to reflect changes that could be expected under hot and dry conditions, similar to the hot and dry scenario developed for the MPRWA.

In a second approach used in Steiner et al. (2000), it was assumed that withdrawal or consumptive use rates for the commercial, industrial, thermoelectric, mining, or livestock sectors would remain constant, but that increases would be seen in the domestic and irrigation sectors. The domestic sector increases were based on the consumptive use rate in 1999, which was a drought year, so the assumption was made that increases in use would be representative of a future hot and dry condition. This rate was multiplied by the forecast number of single family households in the study area for future years. Irrigation sector increases were based on use rates in 1995, which was not a hot or dry year, and then increased by 35 percent across an entire year to simulate conditions in a drought. To forecast for future years, the estimate of consumptive use was applied to the expected future number of irrigated acres.

The results of the Steiner et al. (2000) study indicate a consumptive use rate of 169.1 mgd in June, July, and August for the drainage area above Little Falls, and an increase in consumptive use of approximately 30 mgd between 2000 and 2030. Those estimates, however, exclude the WMA utilities upstream of Little Falls which are included in the scenarios generated in this study. When the WMA utilities are excluded from water use totals in the MPRWA study's future scenarios, then the basin-wide water use totals are similar. At the HUC8 watershed scale there are significant differences between the studies in predicted water use which probably reflects different assumptions and data sources for each water use category (Appendix B, Table 23).

### 4.4 Summary

Total consumptive use above Little Falls across all days in the base year 2005 was estimated to be 385 mgd. Domestic and public supply accounted for 84 percent of this amount. Agriculture, power production, and industry accounted for the remaining 16 percent of consumptive water use. These water use sectors, while accounting for a small fraction of the total water lost through natural and anthropogenic causes on a basin-wide and annualized basis, can be more significant within specific subwatersheds and during extreme low flow events. Scenarios projecting water use to 2030 suggest that domestic and public supply will continue to account for most of the basin-wide consumptive water use. The rate of increase in total consumptive use, if there is no change in the per capita use of water for domestic and public supply, is estimated to be about one mgd per year, which is roughly

consistent with estimates from previous studies. Flow simulations that incorporated future water uses as well as future land uses show these changes will alter multiple flow statistics from their current levels and, given the FA-E relationships identified, these future changes are likely to increase the risk of degraded ecological health.

# Chapter 5. Stream Environmental Flow Needs

---

**Chapter Summary:** The third component of the MPRWA project used the Ecological Limits of Hydrologic Alteration (ELOHA) approach (Poff et al. 2010) to investigate biological responses to flow alteration in streams and small rivers in the Middle Potomac study area. The study component (1) identified flow metrics that, when altered, associate with strong biological responses, and (2) developed flow-alteration ecological response relationships that describe how aquatic ecology responds to flow alteration in a semi-quantitative manner. The results can be used to aid in the future development of environmental flow recommendations for tributary streams. This chapter and six supporting appendices (C-I) describe in detail how the ELOHA framework was applied. Streams and small rivers are defined as Strahler order 4 or less and wadeable; in the study area these waters correspond to the 1a, 1b, 2, and smaller 3 size categories of the Northeast Aquatic Habitat Classification System (NEAHCS). A major part of this MPRWA component involved pairing biological sampling data with observed and/or simulated flow data at numerous, diverse stream locations across the study area. Biological data were extracted from a Chesapeake basin-wide macroinvertebrate database compiled from 23 federal, state and local sources. Observed flow data corresponding to each biological sampling site were limited because the majority of Potomac flow gages are located on rivers. Hydrologic modeling was required to simulate flow time series at ungaged locations. The Chesapeake Bay Program’s Phase 5.2 HSPF model and the Virginia Department of Environmental Quality’s Online Object Oriented Meta-Model (WOOOMM) routing module were used to estimate daily streamflows for 747 watersheds draining to biological monitoring locations. Time series were simulated for baseline and current conditions in each watershed (water years 1984-2005). The current scenarios were validated with observed (gage) data. Five future scenarios were also simulated at 153 of the larger HSPF watershed model segments. The difference between baseline and current scenarios measures the amount of hydrologic alteration already present in a given watershed; the difference between current and future scenarios estimates potential future alteration. A suite of flow metrics was calculated from each scenario time series. The responses of family-level metrics of stream macroinvertebrate to alteration in representative flow metrics were examined with three statistical methods: Pearson correlation, quantile regression and conditional probability. Conditional probability plots were used to create flow alteration-ecology (FA-E) curves. These curves provide a scientific basis for defining environmental flows that sustain healthy stream communities as well as the goods and services that humans derive from well-functioning stream ecosystems. All the data used in this component of the study can be found in Appendix C. Appendix D provides extensive detail on the application of the ELOHA methodology in this study. Appendix E contains technical memoranda describing details of the hydrologic model. Appendix F provides an in-depth comparison of flow metrics calculated from observed (gaged) and modeled time series. Appendix G contains an analysis of nine possible stream classifications (the classification system eventually selected is presented in Appendix D, section 4). Appendix H provides a detailed look at hydrologic alteration in the Middle Potomac study area today and explores possible links to different land uses. Appendix I provides additional details about the FA-E relationships.




## 5.1. Ecological Limits of Hydrologic Alteration (ELOHA)

The ELOHA framework was adapted for the stream and small river environmental flow needs assessment (Figure 4). The four major scientific steps in the ELOHA approach are: (1) build a hydrologic foundation with models that can produce hydrographs for different watershed conditions, including an undisturbed (baseline) watershed, (2) classify rivers according to flow regimes and geomorphic features to reduce biological variability, (3) compute flow alteration using undisturbed hydrological conditions as the baseline, and (4) formulate FA-E response relationships for environmental flows. These four steps were followed in the MPRWA with a little modification. These steps are discussed in detail in Appendix D, Application of ELOHA.

Several important characteristics of the Middle Potomac study area and available data shaped the project and/or required modifications to the prescribed ELOHA process. These included:

- Biological monitoring programs and methods. The Middle Potomac is an interstate watershed, with most of the significant biological monitoring conducted by loosely coordinated but independent state and local agencies using their own methods. A review of the data showed that benthic macroinvertebrates were the only taxonomic group for which comparable metrics could be calculated across jurisdictional boundaries and for which there was a sufficiently large number of samples to allow application of the ELOHA method. Benthic macroinvertebrates are



**Figure 4. Conceptual diagram of the major steps in developing flow alteration-ecology (FA-E) relationships for the Middle Potomac study area.**

This diagram is a framework and the stream/small river component of the MPRWA is one piece of the framework. The black box indicates ongoing monitoring programs that provide the information needed to track changes in flow and stream biota; the green boxes indicate analysis steps taken in the stream/small river component of the MPRWA for the purpose of determining FA-E relationships; the orange boxes indicate both ongoing and future steps in the states' planning efforts, which are supported by this project's outreach and consensus-building efforts. Adapted from Poff et al. (2010).


commonly used indicators of ecosystem health, but this project's reliance on macroinvertebrates, though necessary, has implications for interpretation of the FA-E relationships that result. Fish, for example, may react to certain flow characteristics that macroinvertebrates do not.

- **Stream classification.** Poff et al. (2010) recommend classifying streams by type to reduce natural variability inherent in biological communities. The step is intended to remove variability from, and strengthen confidence in, the FA-E relationships. An established stream classification approach (Olivero and Anderson 2008) was explored in this project; it proved only somewhat successful because it did not account for the confounding influences of other anthropogenic stressors, which can be significant in the Middle Potomac study area. When stream classification was attempted with a subset of the data comprised of the highest quality (reference) sites, with good water quality and habitat conditions, a strong classification system based on the USEPA Level IV ecoregions was found. Ecoregions have distinct regional differences in topography, soils, geography, and vegetation – all of which are factors that affect the structure and function of stream macroinvertebrate communities (e.g., Feminella 2000, Hawkins et al. 2000, Kennan 1999). In these high quality streams, anthropogenic influences are minimized, macroinvertebrate communities are primarily governed by natural features of the ecoregion, and ecoregion differences in the communities are significant ( $p < 0.01$ ). Stream size (Strahler 1<sup>st</sup> – 4<sup>th</sup> order), season (spring, summer, autumn/winter), and karst geology proved less important as classification factors explaining natural variability. A scoring protocol based on each ecoregion's reference communities can be used to evaluate macroinvertebrates anywhere in the basin on a comparable scale of 0% to 100%, thus obviating the need to analyze the data separately by ecoregion classes.
- **Flow metrics.** Although similar studies in other areas have identified FA-E relationships for particular flow characteristics there was no *a priori* reason to believe that the available data in the Potomac watershed would identify those relationships. Therefore, the data were used to define, rather than to test, FA-E relationships. Computer programs for calculating many flow metrics were available and so a large suite of flow metrics and biometrics were calculated. Then, through a series of analytical steps, the list of metrics was winnowed to a very small set that represent different aspects of the flow regime, show a relationship with watershed and water use factors, and show a relationship between flow alteration and biological status.
- **Confounding factors.** It is universally recognized that stream biological status is affected by many natural and anthropogenic factors interacting in complex ways that cannot be easily disaggregated. Figure 5 is a conceptual diagram showing biological community health as the result of multiple processes, of which flow is just one. A challenge of the MPRWA was to distinguish macroinvertebrate changes caused by altered flow from macroinvertebrate changes caused by other environmental factors such as poor water quality and in-stream habitat disturbance. Then, statements could be made about anthropogenic activities that alter flow and the effect of flow alteration on biological community health. As with any data analysis, it is imperative to remember that each dataset has its own particular distribution of natural watershed characteristics and anthropogenic impacts. These features will influence the flow alteration – ecology relationships, either by limiting avenues of investigation or more heavily weighting some results.

Several general premises underlie the ELOHA approach used in this study, namely:

- (1) humans can alter the stream flow regime of a watershed through direct (impoundments, withdrawals, discharges) and indirect (land uses) actions;
- (2) status of the macroinvertebrate community is a function of, among other factors, stream flow regime;
- (3) macroinvertebrate status can be represented by a diverse set of family-level metrics which quantify community composition and function;
- (4) key features of the flow regime (magnitude, duration, frequency, and rate of change) are important to macroinvertebrate communities;
- (5) increasing alteration in these features, expressed as flow metrics, away from a baseline condition corresponds to changes in macroinvertebrate status, generally for the worse; and
- (6) the flow alteration signal can be separated from other, natural and anthropogenic factors affecting macroinvertebrates (e.g., bioregion, pollution).

These premises are derived from the large body of literature on flow effects, which has been discussed and summarized by multiple authors going back to Hynes (1970). The term “premise” is used rather than the recently coined “flow hypothesis” because the project’s analyses do not follow the scientific method and do not test the statements above in controlled experiments. Rather the study’s results and their interpretations lend support to—or counter—the statements by revealing consistent associations between biological communities and conditions with altered flow.



**Figure 5. Conceptual diagram of biological community health.**

Flow is only one of multiple factors affecting biological community health.

## 5.2. Findings for Stream Environmental Flow Needs: Interpretation and Use of Flow Alteration-Ecology Relationships

Three methods were used to examine macroinvertebrate responses to flow alteration: Pearson correlations, quantile regression plots, and conditional probability plots. Each has its strengths and weakness, which are discussed in detail in Appendix D and summarized in Table 6 of Appendix D. Each could be used in specific circumstances to examine macroinvertebrate responses to flow alteration.

The Pearson correlations and quantile relationships demonstrate that alteration in multiple flow metrics has a negative impact on biological health, but also have drawbacks that complicate their interpretation. First, the calculated correlations and regressions were linear. Examination of scatter plots suggests the relationships may not be linear. Second, most macroinvertebrate metric values vary naturally by bioregion and to a lesser extent by season and stream size. Some biometric values can qualify as reference in one bioregion but not in another, which can lead to ambiguous results when the data are pooled without consideration of these natural factors. These limitations could be overcome with (1) a larger dataset collected from a greater diversity of sampling locations within each bioregion, and (2) for each bioregion, an adequately broad range of watersheds experiencing flow alteration. Bioregion-specific FA-E curves could then be developed and used.

The conditional probability method requires a large dataset to produce meaningful results, but it removes much of the underlying natural variability through the use of biometric scores instead of values. Including scores of fair in the calculation reduced additional variability relating to season and stream size in this study (Appendix D, section 4). The use of LOESS regressions to establish the probability curves also removed the linearity imposed on the relationships in the two other methods. As discussed below (section 5.2.6), the conditional probability curves can in theory be shifted by non-flow changes to water quality and in-stream habitat conditions. At this juncture, however, the curves are most suited for developing FA-E relationship curves.

The final suite of macroinvertebrate and flow metrics used to establish FA-E relationships for the Middle Potomac study area are listed here:

### Macroinvertebrate metrics

Chessie BIBI (a multi-metric community index)  
 %EPT (pollution-sensitive group)  
 FBI (Hilsenhoff family-level biotic index)  
 %Chironomidae (pollution-tolerant group)  
 %Scraper (feeding group)  
 %Clinger (habit group)  
 SW (Shannon-Wiener diversity index)

### Flow metrics

3-day maximum  
 High flow index MH21  
 High flow duration DH17  
 Low pulse duration  
 High pulse count  
 Flashiness

Each of the seven biometrics represents an important but different component or function of the macroinvertebrate community; each of the six flow metrics represents a particular section of the hydrograph and has a relationship to land or water uses in the watershed. The metrics are not necessarily the “best” ones for all uses but they serve the purpose of illustrating the array of FA-E relationships in the Middle Potomac study area. The 42 FA-E plots (7 x 6) are presented in

Appendix I. All plots of flow metric percent alteration versus land use can be found in Figure 10 of Appendix H.

Factors that are typically responsible for flow alteration such as dams, withdrawals, and discharges are significant in some ELOHA watersheds but the amount of biological data available to associate with these factors is limited. Urbanization—and impervious surface area in particular—was found in this analysis to be the major factor altering flow regimes in most of the ELOHA watersheds and in the Middle Potomac study area in general. The final FA-E plots display the direction of flow alteration (positive or negative) that has adequate data and exhibits strong flow-macroinvertebrate relationships. Summarized below are the study’s major FA-E findings.

### 5.2.1. Magnitude


All seven macroinvertebrate metrics related strongly to alteration in high magnitude flow metrics but none related significantly to alteration in median flow or any of the low magnitude flow metrics. As a result, FA-E relationships were only developed for high magnitude flows, represented by the IHA 3-day maximum metric. Simulated baseline values of the 3-day maximum are consistently lower than observed reference values, indicating the model underestimates high magnitude flows. However, the metric’s baseline responses to natural factors (watershed size, gradient, karst) and current responses to anthropogenic factors paralleled those in the observed data (Appendix F). Simulated metric values appear to be internally consistent despite their displacement from observed values and were used to develop FA-E relationships.

Results from this study indicate that the 3-day maximum flow metric has increased in more than half of the ELOHA watersheds (i.e., current values higher than baseline values). Positive alteration and percent impervious surface area are very closely related after percent impervious area exceeds a threshold of about one to two percent. Positive alteration in 3-day maximum is also linked to percent urban area, but not as strongly. About a quarter of the ELOHA watersheds showed no alteration (zero percent); the remaining watersheds showed slight negative alteration (current values lower than baseline values) and are often heavily agricultural.

All seven macroinvertebrate metrics exhibit a much lower probability of fair or better status as 3-day maximum increases from baseline (Figure 6). The biometrics %Scrapers, Shannon-Wiener diversity index, and Chessie BIBI show the largest, steepest declines from baseline percentages of fair or better than fair status; %EPT and %Chironomidae show the smallest declines. The biometrics %Clinger, %Scraper, and %Chironomidae drop most quickly as alteration increases from baseline.

### 5.2.2. Duration

All seven macroinvertebrate metrics declined in relation to changes in the duration of both high magnitude and low magnitude flows. Three metrics were used to represent duration: the HIT high flow index MH21 metric, the HIT high flow duration DH17 metric, and the IHA low pulse duration metric. MH21 and DH17 are calculated differently. MH21 is the volume of the high flows (cubic feet) divided by median daily flow (cubic feet per day), where high flows are all those above the study period’s overall median. DH17 is the average duration of events (consecutive days) above the study period’s overall median. Low pulse duration is the median of the annual average number of days per year that flow persists below the study period’s overall 10th percentile. All three metrics are well characterized by the project model.




**Figure 6. FA-E relationships for positive alteration (increase) in annual 3-day maximum flow.**

The Loess smoothed regression lines of the conditional probability plots are shown. “Fair” status of each biological metric is defined by the bioregion-specific thresholds (T%ile) listed in Table 3 in Appendix D.


Negative alteration from baseline indicates flow events have shorter durations. A little over half of ELOHA watersheds exhibit negative alteration in low pulse duration and approximately two-thirds exhibit negative alteration in the two high flow duration metrics. Negative alteration and percent impervious surface area are very closely related after percent impervious area exceeds a threshold of about 0.3 - 0.4 percent. Watersheds with substantially shorter low pulses but relatively unchanged duration of high flow metrics tend to be agricultural. A number of ELOHA watersheds with particularly large discharges no longer have low pulses, meaning their current flows never drop to the lowest levels experienced in the baseline. Many of these same watersheds show a corresponding increase in their high pulse counts. The Eagle Run watershed near Martinsburg, WV no longer experiences any low pulses and its entire simulated current flow time series is above the baseline threshold for high pulses. The watershed is estimated to have discharges equaling 101.5 percent of median flow.<sup>10</sup>

All seven macroinvertebrate metrics exhibit a lower probability of fair or better status as the duration of both high or low flow events decreases from baseline (Figure 7 and 8). The biometrics %Scrapers, %Clingers, and Chessie BIBI show the largest declines from baseline percentages of fair or better than fair status; %EPT again shows the smallest declines.

<sup>10</sup> This watershed was not included in the Pearson correlations because it throws the result off so much.



**Figure 7a-b. FA-E relationships for negative alteration (decrease) in the high flow duration metrics DH17 (A) and MH21 (B).**  
See Figure 6 heading for details.



**Figure 8. FA-E relationships for negative alteration (decrease) in low pulse duration.**  
See Figure 6 for details.

### 5.2.3. Frequency

In the ELOHA watersheds, all seven macroinvertebrate metrics were strongly related to alteration in the frequency of high pulses and to a change in the number of reversals in flow; it is difficult to discern any responses to altered frequencies of low pulses or extreme low flows. The final selection of FA-E plots was for the high pulse count metric. The model appears to have difficulty representing the effect of gradient on number of reversals, which is the reason this flow metric is excluded at this time. High pulse count is the median of the 21 annual averages of high pulse events (consecutive days) in the study period. Flows exceeding the study period's overall 90th percentile are classified as high pulses. Flow metric testing shows the model does an excellent job of representing high pulse count.

Roughly 60 percent of ELOHA watersheds show no alteration in high pulse count. Alteration, when it occurs, is most often in the positive direction, meaning an increase occurs in the number of high pulses. Positive alteration corresponds significantly with percent impervious surface area in the ELOHA dataset when impervious surface exceeds a threshold of about one percent. As percent impervious surface area increases the frequency of high pulses increases. The uncommon instances of negative alteration (fewer pulses) tend to occur in watersheds with large water manipulations such as discharges, withdrawals, and regulated impoundments.


All seven macroinvertebrate metrics exhibit a lower probability of fair or better status when the number of high pulse counts increases from baseline (Figure 9). The biometrics %Scrapers and Chessie BIBI show the largest declines from baseline percentages of fair or better than fair status, followed by FBI and the Shannon-Wiener diversity index. %EPT, %Chironomidae, and %Clinger show the smallest declines in response to more high pulses.

### 5.2.4. Rate of Change


In the ELOHA watersheds, all seven macroinvertebrate metrics were related to alteration in the amount of day-to-day change in daily mean flow. Flashiness, and specifically the Richards-Baker Index (Baker et al. 2004), is the flow metric representing rate of change. The absolute values of all day-to-day changes in daily mean flows are summed for the entire study period and divided by the sum of all the daily mean flows. The higher the index value the flashier the stream. Flashiness is well characterized by the project model.

Roughly three-quarters of ELOHA watersheds exhibit positive alteration from baseline in the flashiness metric. Positive alteration indicates an increase in the metric's value. The metric is most closely related to impervious surface in the ELOHA watersheds. Flashiness increases sharply from baseline when impervious surface exceeds a threshold of about one percent. Agriculture typically has no impact on the metric. Significant withdrawals can enhance positive alteration in flashiness; discharges in some circumstances reduce flashiness (negative alteration).

All seven macroinvertebrate metrics exhibit a much lower probability of fair or better status as flashiness increases from baseline (Figure 10). The biometrics %Scrapers and Chessie BIBI show the largest declines from baseline percentages of fair or better than fair status, followed by %Clinger, %Chironomidae, FBI, and the Shannon-Wiener Index. The %EPT metric showed the weakest response to flashiness.



**Figure 9. FA-E relationships for positive alteration (increase) in high pulse count.**  
See Figure 6 heading for details.



**Figure 10. FA-E relationships for positive alteration (increase) in flashiness.**  
See Figure 6 heading for details.

### 5.2.5. Possible Mechanisms Underlying FA-E Relationships

Annual low magnitude flows were represented in the study by five metrics. They are, in descending order:

August median > Q85Seas  $\approx$  Baseflow Index > 3-day minimum > 1-day minimum

The absence of biological responses to negative alteration (lower magnitude) in the five annual low magnitude flow metrics above, as well as median flow and the inter-annual 7Q10 metric, appears to be real. Of the 1,155 records in the ELOHA dataset, negative alteration occurred in median flow in 43.6 percent of records, in August median in 45.8 percent of records, in the Maryland Q85Seas metric in 31.7 percent of records, in the Baseflow Index in 32.8 percent of records, in the 3-day and 1-day minima in about 38 percent of records, and in the 7Q10 in 14.7 percent of records. Thus, sample sizes were large enough to avoid spurious relationships. Negative alteration in the August median and Baseflow Index flow metrics was significantly related ( $p < 0.01$ ) to the Chessie BIBI, one of the more flow sensitive biological metrics, but the relationships explained little of the biological variation ( $r^2 < 0.05$ ). No significant relationship between negative alteration and the Chessie BIBI was found for median flow or the other low magnitude flow metrics. Similar results were found in the quantile regression analysis. Negative alteration in the median and 3-day minimum metrics did not correlate significantly ( $p < 0.01$ ) with the Chessie BIBI, Shannon-Wiener Index, FBI, %EPT, %Chironomidae, %Scrapers, or %Clingers (Appendix I).

The lack of response by macroinvertebrate metrics to alteration in low magnitude flow metrics suggests their communities, as presently sampled by monitoring programs in the Potomac watershed, are not negatively impacted by existing levels of flow alteration in the magnitudes of normal or seasonal low flows. This result apparently is not an artifact of springtime sampling protocols. Greater than half of the macroinvertebrate data used in the study were sampled in spring and not in the annual low flow period. Spring sampling of macroinvertebrate communities will not necessarily reflect a single low flow event from the previous year; however, flow alteration that results in consistently higher or lower low flows could potentially interfere with summer reproductive timing and success in ways that could affect communities the following spring. This was not observed with the available data.

Recovery from extreme annual low flows may be swift among certain macroinvertebrate taxa that have acquired adaptations to periodic drought events (Humphries and Baldwin 2003). Multivoltine taxa may be able to recover from a seasonal low flow event through successive generations by the time a spring sampling occurs. Highly mobile taxa are also more likely to recover through rapid recolonization from refugia locations (Lake 2003). Additionally, taxa active in the spring may employ strategies that maximize reproductive success by avoiding summer drought impacts such as terrestrial eggs or drought-resistant egg or pupae life stages (Boulton 2003, Lytle and Poff 2004). Taxa that do not display adaptations for rapid dispersal or employ strategies to avoid seasonal drying events may show a lag in recovery. The diverse adaptations of stream macroinvertebrates to periodic dryness make them poor indicators of negative alteration in low magnitude flows.

The 7Q10, or lowest stream flow for seven consecutive days that is expected to occur once in ten years, was used in the study to represent inter-annual low flow magnitudes. Two low flow events typically meet the 7Q10 threshold in a 21-year study period such as this one. Since the study period overlapped the region's drought of 1999-2002, the low levels experienced during this drought would

be reflected in the 7Q10 threshold. Macroinvertebrate metrics did not respond to alteration in the 7Q10. This result was anticipated. First, current and baseline 7Q10 flow levels as depicted by the project's model are approximately the same despite the fact that percent alteration ranges from -100 percent to upwards of 1,000 percent. In baseline scenarios of the ELOHA watersheds, the 7Q10 thresholds were on average 6.2 percent of median flows and ranged from zero percent (no flow) to as high as 26 percent. In current scenarios for the same watersheds, 7Q10 thresholds averaged 6.7 percent of median flows and again ranged from 0 percent to 26 percent with the exception of a few extremely high values associated with high discharges or large impoundments. Anthropogenic impacts for the most part do not appear to alter inter-annual low magnitude flows in the ELOHA dataset. Second, anecdotal evidence from the 1999-2002 drought suggests macroinvertebrates in the Potomac mainstem and larger tributaries were not negatively impacted by the extreme low flow levels (Appendix A). Finally, cross-year comparisons of BIBI scores did not reveal significant decreases in the drought period of 1999-2002 in summer or fall samples.

The combinations of anthropogenic factors present in the ELOHA watersheds tend to raise annual high magnitude flows above baseline levels, sometimes more than doubling peak daily flows. Alterations in annual high flows, represented in the study by the average annual 3-day maximum, inflicted consistent deleterious impacts on all seven of the macroinvertebrate metrics. The literature points to several possible mechanisms for these effects, including scour of periphyton and organic material, catastrophic accidental drift, stranding, alteration of stream-bed habitat, and the indirect effects of poorer water quality (Bunn and Arthington 2002, Poff and Zimmerman 2010, Richardson et al. 2004, Robinson et al. 2004). The particular sensitivity of Scrapers to high magnitude flow supports the concept of periphyton scour as a driver of stress. The strong response of the multi-metric Chessie BIBI coupled with those in the pollution-sensitive metrics, pollution-tolerant metrics, and Shannon-Wiener Index of diversity indicates that overall community structure is compromised by larger annual high magnitude flows. The modest response of the %EPT metric may be due to a lack of sensitivity within the metric. Many taxonomic families and genera comprised the EPT metric. The loss of flow-specialist taxa may result in replacement by more tolerant and flow-generalist taxa, the net effect being little change in a %EPT metric. One such example could be illustrated among caddisflies; *Macrostemum*, a net-spinning caddis fly that constructs firm refugia cemented to benthic substrate would likely be tolerant of peak flows, while other free-living taxa that cling to epibenthic surfaces may be susceptible to dislodgement (Holomuzki and Biggs 2000).

The impacts on macroinvertebrate community status of high and low flow events that are shorter and more frequent, with faster rates of change in daily flow between events, are likely related and share underlying mechanisms. Flow metrics that express duration, frequency, and rate of change in the hydrograph are different in focus but linked by the events that shape them. In urbanized watersheds, the frequency of high flow events and their rates of change increase as hardened surfaces deliver more water more quickly to stream channels, while the duration of flow events decrease. In this study, all seven biometrics showed signs of stress in response to changing frequency, duration, and rate of change in flow. Several underlying mechanisms can be implicated to explain how macroinvertebrate communities, over time, become impacted by the combined features of more frequent and flashier flow events of shorter duration, including: catastrophic drift, displacement from habitat and stranding, and interruption of development or dispersal cues (Lytle and Poff 2004, Poff and Zimmerman 2010, Richardson et al. 2004, Robinson et al. 2004).


The %Scrapers and Chessie BIBI metrics demonstrate the most abrupt declines with alteration in duration, frequency and rate of change flow metrics; %EPT was overall the least sensitive. The

sensitivity of the %Scraper metric can be explained by that group's reliance on periphyton food, which is washed downstream by scouring high flows. Scrapers have been found to demonstrate greater sensitivity to changes in both flow duration and magnitude in other studies (Kennan et al. 2009). In addition to bottom scouring, several other ecological mechanisms can be implicated to explain the flow sensitivity of macroinvertebrate metrics. Rapid changes in flow (flashiness) when combined with unstable bed loads can cause more frequent occurrences of "catastrophic drift" and overwhelm an invertebrate's ability to find refugia resulting in decreases in abundance and richness (Richardson et al. 2004). More frequent high flow events increasingly interfere with longitudinal dispersion triggers, cues for reproduction and development, and the likelihood of coinciding with a flow-susceptible life stage (Poff and Zimmerman 2010). More frequent disturbances do not allow flow-sensitive taxa time to recover and leads to changes in lotic community structure (Poff and Ward 1989).

### 5.2.6. Other Environmental Stressors

Although flow is sometimes considered the master variable (e.g. Poff et al. 2010) and even the primary predictor (Carlisle et al. 2010) of stream biological integrity, it is not the only factor stressing macroinvertebrate populations in the Middle Potomac watershed. As shown in all the quantile regression plots (Appendix I), poor biometric values can and do occur in streams with little or no flow alteration. Further investigation of these poor-performing streams usually uncovers the presence of other stressors such as low pH, high conductivity, low dissolved oxygen, nutrient enrichment, an altered stream channel, poor substrate, or an embedded stream bottom. When streams in the ELOHA dataset with documented high quality are analyzed separately from those with documented degraded quality, the effects of non-flow factors become readily apparent. An example involving the Chessie BIBI and flashiness is shown in Figure 11. Panel A consists of the subset of stream sites with documented high quality habitat and water conditions; panel B has all the ELOHA stream sites; and panel C consists of the subset of stream sites with known habitat and/or water quality problems. The 90<sup>th</sup> percentile regression line (red) trends downward in all groups as flow alteration diverges from baseline, indicating the limits on biological condition set by flow alteration. The 90<sup>th</sup> percentile regression line in the degraded group (C) is positioned lower than in the high quality group (A), reflecting the overall lower macroinvertebrate scores in the degraded group. The line also has a steeper slope. The 90<sup>th</sup> percentile regression line in the middle panel with all the ELOHA data (B) is intermediate to those in the high quality and degraded groups. One can infer from these graphs that if water quality and physical habitat conditions are significantly improved, biological scores will increase regardless of the level of flow alteration at the site. One can also assume that if water quality and habitat conditions remain the same at a site, increasing flow alteration will degrade the biological community.

Conditional probability curves similarly respond to significant water and habitat quality changes. The curves are displaced downward (e.g., there is a lower probability of "fair" or better status) when actions in the watershed increase the overall impact of the non-flow stressors such as pollutants or temperature but do not alter the flow regime. Ameliorating the impacts of non-flow stressors through water quality improvements and stream habitat restoration raises the conditional probability curves.



**Figure 11. Chessie BIBI vs percent alteration in flashiness in high quality streams (A), all streams (B), and degraded streams (C) in the ELOHA dataset.** Red line: 90<sup>th</sup> percentile regression line. Black line: linear regression line. High quality streams (A): habitat scores  $\geq 16$  of 20 and pH  $\geq 6$  and DO  $\geq 5$  mg/L and conductivity  $\leq 300$   $\mu$ S/cm and turbidity  $\leq 50$  NTU. Degraded streams (C): water quality is poor (pH  $< 6$ , DO  $< 5$  mg/L, conductivity  $> 300$   $\mu$ S/cm, or turbidity  $> 50$  NTU) and/or physical habitat score  $\leq 12$  of 20. There are 78 data records in A; 1,155 in B; 51 in C. No evaluation was made of nutrients, toxic pollutants, or other factors that also impact macroinvertebrates. Chessie BIBI scores above 30% (arrow) are “fair” or better.


### 5.2.7. Using the FA-E Curves

If water quality and stream habitat conditions are not expected to change significantly, then the conditional probability relationships established in the preceding sections can be developed into tools for relating a proposed watershed change to a change in flow characteristics and ultimately a likely change in stream health. Figure 12 provides one illustration of how a proposed (future) increase in watershed impervious surface area can be evaluated for its potential impact on stream macroinvertebrate condition. The flow metric in this example is flashiness. The blue arrows in Figure 12A indicate a hypothetical watershed with a current condition of approximately one percent impervious surface area and minimal discharges, withdrawals, and impoundments. The proposed change in the watershed is projected to increase impervious cover to two percent (red arrows). Since the current condition is above the threshold for unaltered flashiness (yellow arrows), there is a strong likelihood that the increase will further alter flow. The red arrows indicate an increase from 10 percent to 20 percent alteration in flashiness can be expected.

If the current and future levels of flashiness are indicated on the corresponding FA-E plots, this information can be translated into an impact on the biological metrics. In Figure 12B, the horizontal axis is percent alteration in flashiness and the vertical axis is the conditional probability of one of the biological metrics, Chessie BIBI, having a fair or better rating. Because many factors affect biological condition, the amount of change in biology cannot be precisely estimated. However, based on the hundreds of samples in the ELOHA dataset, it is apparent that the probability of a Chessie BIBI status of fair or better declines as alteration diverges from baseline. Figure 12C-E demonstrate the relationship between flashiness and %EPT (Figure 12C), %Scrapers (Figure 12D), and %Clingers (Figure 12E).

In some states, numeric limits of flow alteration have been established and applied to FA-E plots relating fish community status to flow alteration (e.g., Michigan, Massachusetts). The identical approach can be applied to this study's macroinvertebrate FA-E plots if agreements on an acceptable level of macroinvertebrate community status—specifically, the probability of a fair or better score—are reached. The choice of measure or measures of macroinvertebrate status used to develop benchmarks can be flexible. However, a suite of macroinvertebrate metrics is recommended in order to outweigh uncertainty caused by the natural and normal variability found in biological communities and to remain sensitive to varying responses to stress among the community constituents. The diverse array of family-level macroinvertebrate metrics tested in this study showed consistent and predictable responses to flow alteration. Further analysis with more specific genus-level metrics should corroborate and perhaps refine these results but probably will not change them.

FA-E curves based on region-specific quantile regressions plots could prove useful to individual states in their 303(d) assessments. Each state in the Middle Potomac study area relies on a suite of macroinvertebrate metrics to decide if stream designated uses are impaired. Some of the family-level macroinvertebrate metrics analyzed in this study are included in these state assessments (e.g., percent EPT, Hilsenhoff FBI). If a sampling site's biometric values indicate impairment according to state thresholds and those values are positioned close to or above the 90<sup>th</sup> percentile regression line when plotted at the stream's current level of flow alteration, chances are good that flow alteration is an important factor impacting the site's macroinvertebrate community.



**Figure 12a-b. An illustration of how watershed characteristics can be linked to flow alteration (A) and flow alteration linked to the Chessie BIBI, a biological response metric (B) in three scenarios: unaltered, current, and future.**

Arrows indicate how values on horizontal (x) axis are associated with values on vertical (y) axis for each scenario. Solid black line (A), log regression relationship between percent alteration in flashiness and percent impervious surface; solid blue line (B), Loess smoothed regression through the conditional probabilities for increments of alteration in flashiness that biological status (Chessie BIBI) is "fair" or better; open red circles, the conditional probabilities; dashed blue lines, 0.05 confidence interval around regression line; solid yellow circle, Chessie BIBI probability of "fair" or better in high quality (reference) streams with no flow alteration.




Figure 12c-d. Additional illustrations of how flow alteration can be linked to biological response for C) %EPT, D) %Scrapers, and E) %Clingers.

### 5.3. Discussion and Conclusions for Stream Environmental Flow Needs

The previous sections describe how the ELOHA framework was applied to streams and small rivers in the Middle Potomac study area for the purpose of building a scientific basis for defining flows that sustain healthy macroinvertebrate communities and the goods and services that humans derive from healthy stream ecosystems. In summary, recursive partitioning analysis of observed (gage) flow data suggested thresholds in various land and water uses that correspond to unaltered or baseline hydrographs. These thresholds were used to model baseline scenarios. Baseline and current scenario hydrographs were simulated for the watersheds of 747 biological sampling locations in the Middle Potomac. This hydrologic foundation yielded a diversity of flow metrics for each watershed and an estimate of alteration from baseline for each flow metric. Comparisons with flow metrics calculated from a Potomac-Susquehanna dataset of gaged flows identified the simulated metrics that most accurately simulated their counterparts in the observed data. Simulated metrics that poorly represented observed metrics and watersheds that did not appear to be accurately modeled were excluded. Anthropogenic causes of flow alteration were examined in the remaining 656 watersheds and FA-E relationships were generated from those watersheds' 1,155 macroinvertebrate samples. A simplified flow chart of the steps used to select the final sample locations, flow metrics, and macroinvertebrate metrics is shown in Figure 13.


Rigorous testing indicates the HSPF watershed model and WOOOMM routing module adequately represent baseline and current scenario flow time series and most of the flow metrics used in this study. Thus, one can reasonably estimate future alteration with flow time series generated with changes in land and water uses representing future conditions. One can also infer the impact of future flow alteration on stream macroinvertebrate status. The impacts of future flow alteration should be an important consideration, given that macroinvertebrate condition is a biocriterion for deciding stream impairment in all Potomac River basin states.

#### 5.3.1. Land Use Change Thresholds of Potential Concern

With the exception of areas underlain by karst geology, the additive impacts of multiple land and water uses on flow appear to be accurately represented by the project's modeling approach (Appendix D, sections 2 and 3). Simulated flows plotted against increasing land use type closely parallel the corresponding graphs of observed flows (e.g., Figure 2 in Appendix F). Flow alteration due to projected changes in land and water uses can be estimated for individual watersheds, and the expected impacts of these changes on stream communities can assist in local and state planning and management activities. Specifically, planners and managers can minimize flow alteration impacts by limiting the amount of impervious surface area, requiring sustainable stormwater management practices, and prioritizing wetland and floodplain restoration and protection.

The plots of both observed and simulated flows suggest that significant alteration in flow metrics from a baseline condition is most often linked to increases in impervious surface and urbanization, and the concomitant loss of forest, in the Middle Potomac study area. In some watersheds, large differences in the quantity of withdrawals relative to discharges, and vice versa, also alter flows. A few large impoundments are located in the study area; however, their operations and subsequent impacts on flow are variable. The impacts of agricultural practices on flow as a whole appear to be comparatively benign. This finding contrast with the well-recognized impacts of agricultural sediment and nutrient runoff on water quality and livestock destruction of unfenced stream habitat.

## Middle Potomac River Watershed Assessment



**Figure 13. Simplified flow chart of the steps used to select the final sample locations, flow metrics, and macroinvertebrate metrics.**

Careful examination of the watershed-to-flow alteration plots for the ELOHA dataset (Figures 10 and 11 in Appendix H) suggests thresholds exist specific to each flow metric and land use. Percent alteration in many flow metrics begins to diverge sharply from zero when land use exceeds 0.4 to 2.1 percent impervious surface or 5 to 15 percent urban land cover or when percent forest falls below 46 to 75 percent (Table 4, all watersheds). Thresholds may also occur in high flow duration metrics (DH17, MH21) when agricultural land cover exceeds 27 percent but other metric responses are weak, variable, and highly confounded by urban land use. Some variability may be due to (a) the fact that the “agriculture” category includes 16 sub-categories ranging from feedlot operations to pasture (Appendix E), and (b) the poor representation of groundwater withdrawals in the model (groundwater withdrawals are incompletely documented). Flow metrics with the lowest (most sensitive) thresholds in each land use type are the durations of high and low flow events and flashiness, which incorporates rise rate, fall rate, and number of reversals. Flow metrics with weak or no obvious thresholds are low magnitude flow metrics such as 3-day minimum, August median, 7Q10, and Q85Seas. These same low magnitude metrics, however, are strongly influenced by direct manipulations of flow volume such as withdrawals, discharges and impoundments. These results

agree to some extent with Brandes et al. (2005) who found that increases in impervious area may not result in measureable reductions in base flow at the watershed scale.

Thresholds apparent in the watershed-to-flow alteration plots in Appendix H are strongly influenced by the mix of land and water uses in each ELOHA watershed. An analysis of selected types of watersheds avoids the additive effects of multiple land/water uses and refines the thresholds (Table 4, only forested and urban watersheds). In watersheds dominated by urban and forest cover and having little or no agriculture or surface water uses (impoundments, withdrawals, discharges), thresholds appear in all of the flow metrics except 7Q10. Increasing impervious cover with the concomitant loss of forest results in flashier streams, with shorter durations of both high and low flows, more reversals, faster rise rates and fall rates, and ultimately more high and low flow events. Thresholds for percent urban cover in these same metrics are approximately an order of magnitude greater than those for percent impervious surface, and flow metric responses to urban land cover are very similar which is not surprising considering imperviousness correlates closely with urban land cover in this group ( $r^2=0.94$ ). For both imperviousness and urban cover, thresholds for flow metrics expressing middle and low annual magnitude flows are relatively high (6.8 – 16.6 percent and 33 – 68 percent, respectively), which suggests urbanization does not strongly affect middle and low flow magnitudes.

If watersheds are jointly dominated by agriculture and forest, and have little or no urban area or surface water uses, clear agriculture thresholds are only found for low magnitude flow and high flow duration metrics (Table 4, only forested and agricultural watersheds). Current levels of the low magnitude flow metrics were higher and durations of the high flow events were longer than baseline levels. This result suggests some water from rain or groundwater is being redirected to surface flows in agricultural landscapes. Differences between baseline (forested) and agricultural watersheds also may reflect lower evapotranspiration rates in pastures and croplands compared to forests. Metrics expressing frequency, rate of change, low flow duration, and inter-annual low flows (7Q10) in agricultural watersheds only intermittently exceed 5 percent alteration from baseline, suggesting other environmental factors are affecting flows. Metrics expressing high and middle magnitude flows in these watersheds show no thresholds. The comparatively benign effects of agricultural land cover on flow found in this study indicates that a loss of forest to agriculture does not currently alter most flow metrics in the Middle Potomac study area.

### 5.3.2. Biological Thresholds of Potential Concern

Macroinvertebrate family-level metrics representing important aspects of community structure and function respond almost immediately when flow changes from the baseline condition. In some biological metrics, change is gradual; in others, change appears to be abrupt. Macroinvertebrate responses were examined with Pearson correlation, quantile regression, and conditional probability. Each method has strengths and weaknesses but all three demonstrate a degrading trend in the macroinvertebrate community metrics when flow changes sufficiently from the baseline condition. Degradation occurs regardless of whether alteration is in the increasing direction (baseline is higher than current) or decreasing direction (baseline is lower than current). Degraded stream communities are less resilient to natural, extreme events and to the multitude of anthropogenic stressors that accompany human activities. They are less capable of processing both natural leaf litter and human

**Table 4. Apparent thresholds of potential concern for different combinations of land uses in the Middle Potomac simulated results.**

Threshold identified when the 20-point moving average of a flow metric’s percent alteration diverges  $\pm 5$  percent from baseline and subsequently continues to diverge. Watersheds with impoundments, withdrawals, or discharges greater than one percent of median annual flow volume are not included in the analysis. Thresholds are rounded to whole numbers, except for percent impervious which is rounded to one decimal place. Key: ? = intermittent change occurs (change may be due to other factors in the watershed) and/or no consistent threshold is found. See text for details, and Appendix H Figure 10 for graphs. \*Possible threshold at ~25-30 percent.

		Magnitude				Duration		Frequency		Rate	
		3-Day Maximum	Annual Median	3-Day Minimum, Q85Seas, Baseflow Index	7Q10	High Flow Duration (DH17), High Flow Index (MH21)	Low Pulse Duration, Extreme Low Flow Duration	High Pulse Count, High Flow Frequency	Low Pulse Count	Rise Rate, Number of Reversals	Flashiness
All watersheds in the ELOHA dataset (n=656)											
%Forest	<	46%	?	?	?	68% - 69%	67% - 75%	46% - 63%	56%	61% - 68%	57%
%Agriculture	>	?	?	?	?	27%	?	?	?	?	?
%Impervious	>	2.1%	?	?	?	0.4%	0.4%	0.5% - 1%	0.7%	0.4% - 0.7%	0.7%
%Urban	>	14%	?	?*	?*	5%	5% - 15%	7% - 12%	10%	7% -	10%
Only forested and agricultural watersheds (<0.35% imperviousness, <1% surface water uses, n=235)											
%Forest	<	?	?	77%	?	65% - 66%	?	?	?	?	?
%Agriculture	>	?	?	20% - 21%	?	30% - 32%	?	?	?	?	?
Only forested and urban watersheds (<5% agriculture, <1% surface water uses, n=152)											
%Forest	<	74%	20%	40% - 45%	?	81% - 82%	81%	77% - 82%	81%	81% - 83%	77%
%Impervious	>	2.0%	16.6%	6.8% - 10.7%	?	0.7% - 0.8%	0.8%	0.7% - 1.6%	0.9%	0.5% - 0.9%	1.6%
%Urban	>	16%	68%	33% - 55%	?	9% - 10%	8% - 10%	9% - 14%	10%	6% - 10%	15%

pollutants. They provide less support to organisms higher in the food web, many of which are economically and recreationally important. In short, they are less capable of providing the goods and services that humans ultimately derive from healthy stream ecosystems.

It is important to ask whether flow alteration or a change in water quality or physical habitat is actually causing the observed biological degradation. As discussed above, many factors influence stream biological communities (Figure 5). Streams with little or no flow alteration can have sampling stations with poor macroinvertebrate status. The quantile regression and conditional probability plots in Appendix I reflect the 2000 – 2009 status quo in Middle Potomac streams with respect to existing levels of nutrients, toxic pollutants, physical habitat damage and alteration, disease, biological manipulation (e.g., fish stocking), and other non-flow impacts. Removing a few of the samples with water quality or habitat issues does not overtly change the quantile regressions, which seems to confirm that a flow alteration impact is present. However, the linear and quantile regressions can shift when stream quality differs considerably (Figure 11). Significant improvements in water quality and physical habitat conditions may counter flow alteration impacts to some extent, and conversely, significant water and habitat degradation may accentuate flow alteration impacts.

One weakness of the quantile regression method is it does not account for bioregion differences in what values of the biological metrics constitute a healthy, or reference, macroinvertebrate community. This weakness is avoided if quantile plots are done separately for each bioregion. Bioregion-specific FA-E curves based on the 90th percentile could then be developed. This approach is difficult to accomplish in the Middle Potomac region because human activities, and hence their flow impacts, differ by bioregion. For example, people prefer to build and farm on flatter land, which in the Middle Potomac is found primarily in the Valleys and Piedmont bioregions. The greatest amount of flow alteration in the Middle Potomac region presently occurs in the Piedmont. At this juncture, only the Piedmont has an adequate breadth of flow alteration and density of data points to begin development of bioregion-specific FA-E curves based on quantile regression plots.

Bioregional differences in the biometric values can also be avoided by scoring macroinvertebrate metrics against bioregion-specific scales such that communities in reference quality stream conditions receive equally high scores. This facilitates direct cross-comparisons of biological status and removes bioregion as an environmental factor causing variability. As shown in Figure 17 in Appendix D, the effect of flow alteration on macroinvertebrate metrics under current conditions can be described by conditional probability curves, with confidence intervals around the curves. The confidence interval reflects spatial and temporal variability in the regression line and is caused by environmental factors unrelated to bioregion. This inherent variability is not static and, like the quantile regression lines, the conditional probability curves respond to significant recovery, remediation, and restoration activities, and conversely, rapid human population growth and urban development. If changes in water quality and physical habitat conditions in the near future are modest, the existing conditional probability curves should hold within their confidence intervals. Analysis of an expanded ELOHA dataset could begin to address the issue of the relative importance of water quality, physical habitat, and flow alteration.

Results of this study suggest it is reasonable to assume that (1) flow alteration will lower any stream's potential to achieve its best possible macroinvertebrate status, and (2) significant changes in other, non-flow stressors will make it difficult to predict exactly how flow alteration will impact biology. On this basis, simulated future scenarios that show flow altering from the baseline scenario or

moving beyond existing or current scenario levels can be expected to degrade the macroinvertebrate community to some extent. The land use thresholds of potential concern in Table 4 can be used directly to determine if flow alteration is occurring or will occur in cases where the watershed meets certain criteria, i.e., forest and urban with little or no agriculture or water uses, forest and agriculture with little or no urbanization or water uses. Models such as the one used in this project or the proposed Computer-Based Evaluation Tool (6.2) may be more appropriate tools for determining the extent of future alteration in stream flow when watersheds have multiple land and water uses.

## 5.4. Technical Information Gaps, Research, and Analysis Needs

Data limitations and technical difficulties encountered while researching stream environmental flow needs for the Middle Potomac region identified information gaps and analysis needs. They are introduced at length elsewhere and summarized here.

### 1. Fish monitoring data collected with consistent, comparable methods

Stream macroinvertebrates, which are typically dominated by relatively short-lived insect life stages, are most responsive to flow metrics that reflect alteration occurring on a regular basis. Their responses to extreme but infrequent events (extreme droughts, extreme floods) occur in the year of the event and then fade as successive generations recover. Fish are longer-lived and may be better indicators of alteration in these extreme flow events and perhaps alteration in summer low flows.

### 2. Groundwater withdrawals

Additional information about groundwater is needed to improve the groundwater component of the HSPF model. Specifically, what aquifer is groundwater removed from and what the impact is of groundwater withdrawals on stream flow.

### 3. Flow metric analysis of a larger dataset of gaged reference watersheds

Comparisons of flow metrics from gaged reference watersheds and simulated baseline conditions build confidence that the hydrologic simulations are accurately representing the natural effects on flow of watershed size, gradient, geology, etc. Presently, there are only 11 reference watersheds in the Potomac-Susquehanna gage dataset. The analysis should be expanded to include other Pennsylvania, West Virginia, and Virginia river basins adjacent to the Middle Potomac.

### 4. Flow metric analysis of more pairs of simulated and gaged watersheds

Direct comparisons of gaged and simulated flow metrics build confidence that the HSPF model coupled with the WOOOMM routing module is accurately representing the multiple anthropogenic impacts on flow in diverse watersheds. Presently, flow metrics for only 53 watersheds have been directly compared. More gaged watersheds should be simulated for the purpose of direct pair-wise comparisons.

### 5. Flow metric analysis of small (<10 mi<sup>2</sup>) watersheds

Small watersheds are under-represented in the gage data, and are of most interest to local jurisdictions. The search for small gaged watersheds should be expanded to river basins adjacent to the Potomac. Comparisons of flow metrics from gaged and simulated small

watersheds build confidence that the HSPF model coupled with the WOOOMM routing module is accurately representing the smallest watersheds.

6. Coastal Plain watersheds

The Coastal Plain physiographic region has experienced, and will continue to experience, significant impacts from agriculture and urbanization. It is also under threat of sea level rise. The Potomac River basin has a few Coastal Plain watersheds adjacent to its estuary. Determining flow alteration impacts on these watersheds would be best accomplished in the context of the larger, Mid-Atlantic region.

7. Karst watersheds

Karst geology underlies a broad swath of the Middle Potomac region which is expected to see some of the largest increases in urban development in the next 20 years. The simulated flows indicate that these watersheds behave differently than the rest in the Middle Potomac; hence, they show up as “oddballs.” It is not known whether this is a good representation of reality or the consequence of a lack of observed data in karst watersheds for comparison. More observed data is needed to determine model efficiency in these karst regions.

8. Relative importance of water quality, habitat condition, and flow alteration in each bioregion

Analysis of an expanded ELOHA dataset could begin to address the issue of the relative importance of water quality, physical habitat, and flow alteration. The addition of watersheds with known flow alteration but documented good water quality would facilitate the analysis, especially in the Ridges and Valleys bioregions.

9. Enhanced groundwater model component

The HSPF model needs to be enhanced to allow it to simulate impacts of groundwater withdrawals on surface stream flow.

10. Impoundments

The Potomac basin is a relatively unregulated system, meaning there are relatively few impoundments in the basin. Sixteen impoundments were simulated as part of this study, including twelve that were not previously simulated in the Chesapeake Bay Program’s HSPF model. This small sample size is problematic, however, in determining the impacts of impoundments on the flow regime. Additional efforts to enhance this understanding could include incorporating additional, smaller dams in to the hydrologic model and/or selecting additional watersheds that are influenced by the simulated impoundments to increase the sample size.

# Chapter 6. Stakeholder Outreach and Implementation Options

---

**Chapter Summary:** The fourth and fifth components of the MPRWA project were, respectively, stakeholder participation and outreach, and development of implementation options. Stakeholder participation and outreach included construction of a website, a series of webinars, two in-person workshops, and Technical Advisory Group (TAG) meetings. Copies in PDF format of the webinar presentations, the streams and small rivers workshop materials, presentations, notes, and summary, and the TAG meeting materials and presentations are available in Appendix J. The large river workshop summary is included in Appendix A. Development of implementation options consisted of a literature review of Decision Support Tools from across the United States and a concept paper that scopes out a basin-wide comprehensive plan for the Potomac River basin. These documents can be found in Appendix K.



## 6.1. Moving the Science into Management

The MPRWA scientific process yielded a quantitative understanding of the biological effects of hydrologic alteration, and associated these effects to watershed drivers such as land and water uses. Water resources decision-makers can utilize this information to manage watershed conditions to achieve desired hydrologic or biologic results. Case studies for utilizing ELOHA results for management purposes in Michigan and Massachusetts (Reeves et al. in review, Weiskel et al. 2010) were presented in the October 2011 webinar (Science to Management Applications, Appendix J). A list of case studies also is maintained by TNC.<sup>11</sup>

Management efforts, however, are dependent on governance drivers such as institutions, regulations, and policies, which often differ between political jurisdictions, making homogeneous basin-wide implementation difficult. To this end, a participatory process has been ongoing throughout the project to encourage communication with stakeholders across the basin. A description of these efforts is included below. Two products have been identified that may be useful in the implementation of the project findings -- a computer-based evaluation tool and a basin-wide comprehensive plan. The quantitative FA-E response curves can be implemented in a computer-based evaluation tool to assist managers in understanding the implications of various land and water management decisions. The results can also be utilized, along with other existing information, for planning purposes in a Potomac Basin Comprehensive Water Resources Plan. Development of these two products is outside the scope of the Middle Potomac River Watershed Assessment; however, preliminary evaluation and scoping were conducted.

---

<sup>11</sup> <http://conserveonline.org/workspaces/eloha/documents/template-kyle>

## 6.2. Stakeholder Participation

To encourage implementation of the MPRWA findings, managers and stakeholders have been included throughout the project. These efforts (documented in detail in Appendix J) include a website, a series of webinars, two in-person workshops, Technical Advisory Group (TAG) meetings, and correspondence with state and federal agencies. The outreach efforts were designed to inform stakeholders of the complex technical work as it was conducted. The benefits of stakeholder participation include enhanced understanding of the results and implications of the study, opportunities for feedback on the technical work, and modification of the study design to ensure development of the most appropriate tools and outcomes for implementation.

The project website<sup>12</sup> was designed to convey information to stakeholders and the general public regarding the MPRWA. Up-to-date information such as the webinar slides, draft reports, and workshop announcements were posted to the website to communicate project activities and outcomes.

The webinars allowed the technical team the opportunity to present methodologies and preliminary findings as well as incorporate stakeholder suggestions on the scientific analyses. The first webinar was held in September 2009 to introduce stakeholders to the project. Subsequently, a multi-part webinar series was conducted to describe the technical approach and associated results. The webinar topics for this series were: (1) overview of Potomac environmental flows project goals, outcomes, benefits, and audiences, (2) human uses of water: current and future demands and impacts on flows, (3) modeling stream flows: explanation of models and scenarios, and descriptive statistics for simulated flow scenarios, (4) developing FA-E response relationships part 1: data, variables, and methodology, (5) developing flow alteration-ecological response relationships part 2: what the relationships look like, dealing with data ambiguities, and interpretation, (6) from science to management implications, and (7) future scenarios. A final project wrap-up webinar was held in June 2012. The presentation portion of each webinar was approximately one hour followed by up to half an hour of discussion. The webinars were an integral part in building an understanding of the project so that the project team could obtain informed feedback. Copies of the webinars (PDF format) are available in Appendix J.

Two workshops were hosted in association with the MPRWA. The first workshop, held in September 2010, was the Potomac Basin Large River Environmental Flow Needs workshop. The second workshop, held in November 2011, was the Potomac Watershed Small Stream Environmental Flow Needs workshop. Both of these events convened a group of experts to review and provide feedback on the small and large river studies, utilizing the ESWM and ELOHA frameworks, respectively. These workshops provided an excellent forum for dialogue on the project methodologies, results, and conclusions. At the streams and small rivers workshop, for example, the technical team received and incorporated feedback on the types of flow metrics that are the most useful for management purposes and methods for conveying the findings in a meaningful way for managers. Participants at the large rivers workshop reviewed draft recommendations and their input is reflected in the final recommendations for Large River Environmental Flow Needs (Chapter 3). Complete compilations of feedback received from participants at the September 2010 large river and

---

<sup>12</sup> [www.potomacriver.org/projects/middle-pot-assess](http://www.potomacriver.org/projects/middle-pot-assess)

November 2011 small stream flow needs workshops are provided as summary reports in Appendix A and J, respectively.

The TAG was established to provide critical feedback on the development of the flow alteration ecological response relationships. It was composed of experts in stream ecology, environmental flows, and watershed modeling. An effort was made to have state-level representation from all basin jurisdictions. The TAG met twice at ICPRB in Rockville, MD, in September and October 2011, leading up to the November workshop. The meeting materials are provided in Appendix J. All TAG members were invited to attend the November 2011 project workshop, where their expertise and knowledge of the project proved valuable in communicating with other stakeholders and providing thoughtful comments on the project.

Early in the study letters were sent to multiple federal and state agencies to convey initiation of the study and request their participation. All basin states and several federal agencies participated in the study, to varying degrees of involvement. Responses to the initial study correspondence were received from the Maryland Department of the Environment (MDE), MDDNR, Pennsylvania Department of Environmental Protection (PADEP), VADEQ, Natural Resources Conservation Service (NRCS), and USGS. Copies of correspondence are included in Appendix J.

### **6.3. Computer-Based Evaluation Tool**

A computer-based evaluation tool could assist water resources managers in assessing potential impacts from development by providing the predicted hydrologic and biologic impacts of a particular management decision. The optimum configuration of a Potomac computer-based evaluation tool is still to be determined; however, the general concept is as follows. The FA-E responses developed as a result of this study provide managers with an understanding of the degree of biological alteration associated with a particular amount of hydrologic alteration. Computer-based evaluation tools can be utilized to look at different “what-if” scenarios to anticipate the degree of biological degradation or improvement associated with a particular activity. Then an informed management decision can be made about proceeding with the activity. Useful applications of this type of tool may include, but are not limited to, local land use planning and water withdrawal/discharge permitting.

Some tools, for different parts of the United States and Canada, have been developed already for similar purposes. A literature review was conducted to obtain additional information about existing tools and to assess the technologies available and costs associated with creating a user-friendly tool useful in basin-wide water resources management (Appendix K). Available tools varied widely in cost and scope. Some tools included web-based spatially explicit mapping capabilities while others were comprised of downloadable software applications. Depending on the application, tools were developed with a focus on the management of groundwater, surface water, or both. The cost of these tools, for organizations reporting costs, ranged from approximately \$100,000 to almost \$2,000,000. The more sophisticated tools often include online interactive capabilities, integration with existing hydrologic models, and various spatial resolutions for analysis.

## 6.4. Potomac Basin Comprehensive Water Resources Plan

The MPRWA included development of a concept for a Potomac Basin Comprehensive Water Resources Plan. Water resources planning efforts have been underway within the jurisdictions of the Potomac basin for some time. Maryland requires that local governments develop Water Resources Elements within their respective comprehensive plans under Maryland House Bill 1141 of 2006. As part of Virginia's Local and Regional Water Supply Planning Regulation of 2005, development of local or regional water plans is required. Pennsylvania has a State Water Plan process that includes the designation of Critical Water Planning Areas under Act 220 of 2002. West Virginia is working towards the development of a State Water Plan as part of the Water Resources Protection and Management Act, scheduled for completion in 2013. The purpose of a Potomac comprehensive plan is to build on these and other efforts to identify and address water resources issues of interstate or basin-wide concern.

The study developed a concept for how to achieve an adaptive basin-wide comprehensive water resources plan to serve as a roadmap for the sustainable use of this interstate resource now and into the future. A comprehensive plan facilitates proactive, integrated management of the water resources in the Potomac basin. Development of the plan itself was outside of the scope of the MPRWA and will proceed as funding sources are identified and become available. It is anticipated that this effort would be led by ICPRB. However, the strategy for developing a basin-wide plan was prepared under the MPRWA (Appendix K).

In general, the planning process should be collaborative, adaptive, integrated, and participatory. Collaboration will be essential to developing a basin-wide comprehensive plan. Working together, a shared vision can be developed and common goals can be met. The plan should also be adaptive. That is, recommendations can be developed based on existing knowledge, re-evaluated after implementation, and revised as necessary to ensure goals are met. The planning effort should include an integrated perspective of water resources management, taking into account not only water quality, but water quantity, not only hydrologic, but biological concerns. And the planning process should be participatory to engage stakeholders from across the basin, ensuring all voices are heard and incorporated.

Because the specifics of the plan would be determined with significant stakeholder feedback, the planning framework is designed in four phases with the intent of being flexible, in anticipation of future adjustments. The four phases of the plan's development are scoping, identification of water resources issues, identification of solutions, and development of the plan. The scoping process was initiated as part of the MPRWA and includes defining roles of the participating organizations, establishing an interdisciplinary oversight committee, and defining the goals of the plan. Identification of water resources issues has been performed at various scales in numerous previous studies in the Potomac basin. The plan would build on the results of the MPRWA and other studies to identify water resources issues of interstate or basin-wide significance. Issues may include, but are not limited to, water availability, water quality, water use, potential climate change, stormwater and impervious cover, source water protections, and flood and drought management. Management alternatives will then be developed and evaluated to address the identified issues, leading to the recommendation of practical solutions. After compiling the results of the planning process, a Potomac Basin Comprehensive Water Resources Plan document will be prepared.

Benefits of the comprehensive plan would be numerous and include, but are not limited to (1) integration of existing data and research for analysis at the basin-wide scale, (2) enhancing interstate collaboration, (3) management the interstate resource of the Potomac basin at the interstate scale, and (4) increasing cost efficiency through collaboration.

Initial stakeholder outreach for the comprehensive plan was conducted as part of the Middle Potomac River Watershed Assessment at an ICPRB Commission meeting in December 2010, a webinar for key stakeholders and Commissioners in March 2011, personal conversations with representatives from each basin jurisdiction, and distribution of the concept document (Appendix K) to key basin stakeholders for review and comment. The comprehensive plan concept was also presented and published in the proceedings of the American Water Resources Association (AWRA) 2011 summer specialty conference on Integrated Water Resources Management: The Emperor's New Clothes or Indispensable Process (held June 27-29 in Snowbird, Utah). Participation in this conference provided the opportunity to receive feedback on the plan from practitioners. A challenge for successful completion of the basin-wide plan noted at the conference will be the difficulty in getting commitment of resources from organizations whose purview only covers a portion of the basin, because these organizations may feel limited responsibility for the entire watershed. Suggestions received from the group included utilizing Memoranda of Understanding to encourage implementation from basin jurisdictions as ICPRB is a non-regulatory entity.

### **6.5. Discussion**

Ultimately, any land or water management decision-making that can be influenced by the findings of the MPRWA and the Potomac Basin Comprehensive Water Resources Plan will be conducted by local, regional, state, and federal managers throughout the basin. The participatory process of this project was intended to inform basin water managers and other stakeholders of the concepts and methods used to evaluate the impact of land use change and water use on stream flows and the consequent impacts on aquatic biology, and to obtain feedback from those experts that helped with interpretation of analytical results. Implementation of the Comprehensive Water Resources Plan is a logical next step. It can incorporate the technical findings of this project in a stakeholder led process to develop regionally appropriate, sustainable, water management. Development of a computer-based evaluation tool, by a project partner, could facilitate understanding of the implications of various land and water use management decisions. USACE could undertake development of a tool through the Interagency and International Support Technical Assistance Program.

# Chapter 7. Conclusions

---

**Chapter Summary:** This chapter summarizes the overarching findings of the study. It also lists all the products of the MPRWA study, lessons learned, information needs with respect to research, monitoring and modeling, some potential applications by the jurisdictions, and next steps in establishing environmental flows for the Middle Potomac River region.



## 7.1. Introduction

The Middle Potomac study area presented both special opportunities and special challenges in this assessment. The assessment was an opportunity to evaluate the effects of flow alteration on stream and river ecology in a watershed with essentially unregulated flow. In this watershed, the causes of flow alteration are dominated not by dams but by land use changes, withdrawals, and discharges. There are opportunities to establish ecological benchmarks of flow alteration because the area has a rich and long history of monitoring both flow and biology in and near flowing waters. Potential impacts of future flow alteration were also investigated since water supply and population growth are of interest to, and studied by, several state and regional agencies. As an interstate watershed overlapping the District of Columbia and parts of Maryland, Pennsylvania, Virginia, and West Virginia, there were special challenges in acquiring comparable datasets. In scoping an approach to develop a comprehensive basin-wide water resources plan, the lack of a single basin-wide regulatory authority was a key driver in shaping the proposed participatory and multi-jurisdictional planning framework.

The Middle Potomac is exceptional in that relatively few large dams regulate water flow in the watershed. The combined storage capacity of all major impoundments in the basin upstream of Washington, DC presently makes up less than seven percent of median annual flow, and most impoundments are run-of-river. Extensive farming and timber harvesting historically impacted streams and rivers, but a majority of subwatersheds have since reforested. Many streams and rivers, particularly in the relatively undeveloped western portion of the watershed, are now considered relatively natural with respect to flow. Population growth and urbanization are larger causes of hydrologic alteration than impoundments. The eastern side of the watershed near Washington, DC has experienced significant urban growth since the 1930s. Population is expected to continue to grow and expand westward in the next decades, increasing water use and consumption and converting more forest and agricultural lands to suburban and urban land.

## 7.2. Study Findings

The MPRWA resulted in a number of findings about the study area, some of which have implications for other studies. These findings include the effects of a naturally highly variable flow regime, drivers of flow alteration in the Middle Potomac, recommendations from project stakeholders, and insights on using the ELOHA framework.

### **7.2.1. The Potomac Large River Environmental Flow Needs Assessment**

The Potomac Large River Environmental Flow Needs Assessment found no current identifiable adverse impact on key biotic taxa due to human modification of flow, but the assessment also identified significant data gaps. Biological data collected in comparable ways in the largest river segments of the Potomac are presently insufficient for flow needs assessments. Independent development of FA-E relationships, such as those for macroinvertebrates in streams and small rivers, is not possible at this time. Furthermore, the FA-E relationships identified in streams cannot be extrapolated to the large rivers due to natural changes found in macroinvertebrate communities as rivers broaden and deepen (e.g., Vannote et al. 1980). Relationships between land use change and change in flow characteristics in small watersheds may be applicable to the larger watersheds. These relationships were consistent across a wide range of drainage areas.

The project team, with expert workshop participants' input, arrived at nine conclusions—the main one being that current large river flow characteristics should be maintained (Section 3.4). In particular, the existing minimum flow recommendations for Little Falls and Great Falls should be maintained until (and if) future ecological monitoring identifies impacts that suggest a need to modify flow management policies. The lack of observed adverse impact may be due to the fact that the Potomac watershed has naturally highly variable flows and the taxa present in the river ecosystems are adapted to the flow regime as it exists today.

### **7.2.2. Future Water Use Projections**

Total consumptive use above Little Falls across all days in the base year 2005 was estimated to be 385 mgd. Domestic and public supply accounted for 84 percent of this amount. Agriculture, power production, and industry accounted for the remaining 16 percent of consumptive water use. These water use sectors, while accounting for a small fraction of the total water lost through natural and anthropogenic causes on a basin-wide and annualized basis, can be more significant within specific subwatersheds and during extreme low flow events. Scenarios projecting water use to 2030 suggest that domestic and public supply will continue to account for most of the basin-wide consumptive water use. The rate of increase in total consumptive use, if there is no change in the per capita use of water for domestic and public supply, is estimated to be about one mgd per year, which is roughly consistent with estimates from previous studies. Flow simulations that incorporated future water uses as well as future land uses show these changes will alter multiple flow statistics from their current levels and, given the FA-E relationships identified, these future changes are likely to increase the risk of degraded ecological health.

### **7.2.3. Stream and Small Rivers Environmental Flow Needs Assessment**

The stream and small rivers environmental flow needs assessment evaluated the impact of watershed characteristics, change in land use/land cover, and human uses of water on ecosystem response. Table 5 summarizes the assessment's flow alteration - macroinvertebrate results and lists the possible underlying mechanisms identified in other studies and discussed in Chapter 5.

**Table 5. Summary of flow alteration – macroinvertebrate relationships identified.**

Possible underlying mechanisms that have been suggested in the literature are also listed.

<b>Alteration in these aspects of the flow regime is associated with:</b>	<b>Degradation in these macroinvertebrate metrics:</b>	<b>Possible mechanisms that could explain the association are:</b>
<ul style="list-style-type: none"> <li>• Higher maximum flows</li> <li>• Shorter duration of high flows</li> <li>• Shorter duration of low flows</li> <li>• More low flow pulses</li> <li>• More high flow pulses</li> <li>• Faster rates of change in flow (flashier)</li> </ul>	<p>All 19 family-level macroinvertebrate metrics and the Chessie BIBI multi-metric index</p>	<ul style="list-style-type: none"> <li>• Scour of periphyton and organic matter (food) during high flows</li> <li>• Catastrophic accidental drift during floods</li> <li>• Displacement from habitat and stranding when waters recede</li> <li>• Physical alteration of stream bed habitat</li> <li>• Indirect effects of poor runoff water quality (sedimentation, pollutants)</li> <li>• Interruption of development or dispersal cues</li> </ul>
<ul style="list-style-type: none"> <li>• Lower middle and low magnitude flows, includes median flow, August median flow, summer Q85 flow, baseflow index, 3-day and 1-day annual minima, and 7Q10</li> </ul>	<p>None of the biometrics</p>	<ul style="list-style-type: none"> <li>• Swift recovery due to adaptations to low flow (drought resistant or diapausing life stages)</li> <li>• Multi-voltine (short) life cycles</li> <li>• High mobility, able to find refugia and later recolonize</li> </ul>

Findings from the stream and small rivers environmental flow needs assessment also included:

**Flow Alteration**

- The plots of both observed and simulated flows suggest that significant alteration in flow metrics from a baseline condition is most often linked to increases in impervious surface and urbanization, and the concomitant loss of forest, in the Middle Potomac study area.
- In some watersheds, large differences in the quantity of withdrawals relative to discharges, and vice versa, also alter flows.
- A few large impoundments are located in the study area; however, their operations and subsequent impacts on flow are variable.
- The impacts of agricultural practices on flow as a whole appear to be comparatively benign.

**Land Use**

- The amount of impervious surface appears to be the largest driver of flow alteration in the Middle Potomac. The second largest driver of flow alteration is disproportionate amounts of withdrawals and discharges (i.e. watersheds with large withdrawals but no associated discharge or vice versa). Agricultural impacts on the flow regime are variable but generally weak, and are often confounded by water uses in other sectors and urban land use.

- Careful examination of the watershed-to-flow alteration plots for the ELOHA dataset suggests thresholds exist specific to each flow metric and land use. Percent alteration in many flow metrics begins to diverge sharply from zero when land use exceeds 0.4 to 2.1 percent impervious surface, or 5 to 15 percent urban land cover, or when percent forest falls below 46 to 75 percent.
- Thresholds may also occur in high flow duration metrics (DH17, MH21) when agricultural land cover exceeds 27 percent.

### Biological Metrics

- Biological metrics were strongly correlated with flow metrics representing high magnitude flows, the duration of high and low flow events, the frequency of high and low flow events, and the rate of change in flow. They were not correlated with alteration in the magnitudes of mid-range and low flows.
- Macroinvertebrate family-level metrics representing important aspects of community structure and function respond almost immediately when flow changes from the baseline condition. In some biological metrics, change is gradual; in others, change appears to be abrupt. The likelihood of a macroinvertebrate community having an acceptable status decreases as alteration increases in most flow metrics
- It is reasonable to assume that flow alteration will lower any stream's potential to achieve its best possible macroinvertebrate status, and significant changes in other, non-flow stressors will make it difficult to predict exactly how flow alteration will impact biology.
- For those watersheds least impacted by other non-flow factors there is a significant decrease in biological condition as flow alteration increases.
- To accurately forecast change in the biological community caused by a specific change in flow projections must account for all the natural and anthropogenic factors in addition to flow alteration that impact biological communities. Degrading water quality and damaged in-stream physical habitat—or conversely water quality improvements and stream rehabilitation—can significantly influence the local biological community.

In addition to estimating hydrologic alteration from baseline to current conditions, alteration was estimated in the Middle Potomac watershed under five future scenarios. Future scenarios that incorporate meteorological changes, including the hot and dry and climate change scenarios, display the most extreme hydrologic alteration throughout the study area because decreases in precipitation and increases in temperature reduce the amount of water available for both human and ecosystem use. In future scenarios that include only changes in land and water uses (DP1, DP2, and power), alteration is primarily associated with decreases in forest and increases in impervious cover (driven by increases in population); although total alteration is a combination of various factors that may work to either temper or exaggerate the overall alteration in a particular location. The largest current and future withdrawals with respect to volume are located in the North Branch and on the mainstem Potomac River, altering the hydrology in the mainstem system. These withdrawals are typically associated with decreases in median flows and flashiness.

An initial goal of the report was to define general, transferable FA-E relationships that represent the “environmental flow needs” of streams and small rivers in the Middle Potomac study area, but the report stops short of achieving this. The quantile regression plots illustrate that macroinvertebrate communities in the study area are locally impacted by factors other than flow alteration, such as poor water quality and disturbed stream habitat (e.g., Figures 21 and 28). The shapes and positions

of the FA-E relationships depend heavily on the proportional impacts of *all* anthropogenic factors in the study area. For example, the response trajectory of biological change in a low quality site stream will roughly parallel that in a high quality stream, but biometric values will be displaced downward because of the additional impacts of poorer water quality and habitat conditions. As Poff and Zimmerman (2009) have concluded, it is not possible to develop “general, transferable quantitative relationships between flow alteration and ecological response” even though flow alteration increasing away from a baseline state is consistently and strongly associated with declining ecological condition.

#### 7.2.4. Stakeholder Engagement Process

The stakeholder process utilized during the MPRWA yielded valuable feedback/insights regarding the project findings. These lessons include a list of flow and biometrics that are of greatest interest for management purposes, methods for presentation of results to inform management efforts, and interpretation of results from the stakeholders’ perspectives. A complete compilation of stakeholder feedback is available in Appendix J.

Defining limits of acceptable hydrologic alteration towards management of ecological health, a final step in the ELOHA process, is a stakeholder driven social or political decision-making process. Limits of acceptable alteration were not defined in this study. Ecological health is a function of many variables, of which flow is only one. Ecological health can be improved by reducing water pollution, improving in-stream habitat, etc. in the absence of restored, naturally variable flow regimes. Acid mine drainage remediation, point source pollution reductions, and agricultural and stormwater best management practices provide examples of this. And, as is often the case in urban areas, flow alteration is sometimes linked too closely with pollutant loads to easily distinguish it as the predominant stressor of the aquatic community. Defining a limit of hydrologic alteration alone to manage ecological health, therefore, is a difficult proposition due to the other confounding factors. This would be an appropriate next step based on the results of this study.

Land and water uses are in many cases subject to local government control. Therefore, actions such as stormwater management that maintain or restore environmentally sustainable flows may need to be implemented at that level of government. Urban stormwater management is capable of reducing peak stormwater flows and slowing the delivery of rainwater to streams and rivers. Actions that address stormwater runoff will reduce the tendency of impervious surfaces in urban areas to increase the magnitudes and shorten the durations of high flows in streams after storm events. Hydrologic alteration associated with disproportionate amounts of withdrawals and discharges may illustrate the hydrologic impacts of transfers between subwatersheds and, in some cases, may be due to limitations in the withdrawal and discharge datasets available for this study.

#### 7.2.5 Summary

In the large rivers included in this study, based on currently available information, there has been no discernible adverse ecological impact on focal species due to human modification of flows. As a precautionary measure, the team recommended that the current large river flow regime be maintained for the entire range of flows as defined by 20 flow statistics based on a 21-year period of record (1984 – 2005). In streams and small rivers, strong relationships were found between urbanization (impervious surface), hydrologic alteration, and ecological impacts. Land use was found to be a more significant cause of hydrologic alteration than water withdrawals and

impoundments at the present time. The impact on flow of future changes in water use showed no regional pattern of alteration that applied to all future scenarios – impacts on flow varied for each subwatershed’s unique combination of land and water uses.

Maintaining healthy streams and watersheds, as reflected through the status of stream biological communities, is important to meet Potomac River and Chesapeake Bay restoration goals and is the vision. Defining limits of acceptable hydrologic alteration towards management of ecological health were not defined in this study. The process of defining societal values and needs, determining desired ecological conditions for specific streams and watersheds, and setting limits to flow alteration can best be pursued at a watershed jurisdictional scale, in accordance with state- and local-level priorities, needs, and regulatory mandates. At the local level, the results may be most applicable for land use planning and zoning. At the state level, the results could influence the development or refinement of state stormwater laws to advance Bay restoration goals, or support the development of new flow-related water quality standards.

### **7.3. Study Products**

The assessment generated datasets and analyses that may be useful in future studies and management actions in the Potomac watershed and elsewhere. Products of the study’s five distinct components are listed in Table 6.

**Table 6. Summary of major products.**

Component	Product Potential Use	Reference
Large River Environmental Flow Needs Assessment	<p>Review of stream flow requirements of fishes, mussels, and plant species in the non-tidal and tidal river.</p> <p>Flow needs synthesis including conclusions about Potomac large river flow needs, identifying specific flow regime component needs for key species.</p> <p>Review by workshop participants with expertise in large river ecosystems.</p> <p>Recommendations to address environmental flow needs information gaps.</p> <p>On-line bibliographic database of literature relevant to environmental flow requirements of the Potomac River.</p> <ul style="list-style-type: none"> <li>• Development of hydroecological monitoring plan</li> <li>• Development of quantitative flow recommendation for mainstem Potomac River</li> <li>• Re-evaluate 300/100 mgd flow-by recommendation</li> </ul>	Chapter 2 and 3 Appendix A
Future Water Use Projections	<p>Estimates of 2005 water withdrawals and consumptive use by economic sector.</p> <p>Estimates of 2030 water withdrawals and consumptive use by economic sector for five different future scenarios.</p> <ul style="list-style-type: none"> <li>• Inform water allocation decisions</li> <li>• Inform water withdrawal/permit decisions</li> <li>• Inform land use decisions</li> </ul>	Chapter 4 Appendix B Appendix C
Stream Environmental Flow Needs Assessment	<p>Datasets for impoundments, withdrawals, discharges, land cover, gaged streamflow, benthic macroinvertebrates, simulated streamflow time series, calculated flow metrics for observed and simulated flow time series, and GIS shapefiles for modeled watersheds.</p> <p>Hydrologic model (HSPF) of Middle Potomac ELOHA watersheds and calculation of simulated and observed flow metrics.</p> <p>Statistical analysis of correspondence between watershed characteristics and flow metrics.</p>	Chapter 5 Appendix D Appendix C  Appendices E-F  Appendices F-G

Middle Potomac River Watershed Assessment

Component	Product Potential Use	Reference
	<p>Flow alteration correspondence with human water use and land use change.</p> <p>Flow alteration – ecology (FA-E) relationships.</p> <ul style="list-style-type: none"> <li>• Development of hydroecological monitoring plan</li> <li>• Develop and implement management efforts associated with land use planning, withdrawal/discharge permitting, or other types of water resource management activities.</li> <li>• Thresholds in the watershed-to-flow alteration relationships can inform managers as to when increases in specific land uses are likely to begin altering stream flows.</li> <li>• Relationships between flow alteration and biological metrics can generally indicate how much flow alteration will impact biological communities.</li> <li>• Explore and determine vectors of ecological degradation in basin rivers and streams – is flow the master variable driving ecological conditions.</li> </ul>	<p>Appendix H</p> <p>Appendix I</p>
Stakeholder Engagement Process	<p>Workshop participant suggestions and comments for Large River Environmental Flow Needs.</p> <p>Webinars</p> <p>Workshop participant suggestions and comments for environmental flows.</p> <p>Correspondence with agencies</p> <ul style="list-style-type: none"> <li>• Build consensus on acceptable levels of biological degradation due to changes in flow regime</li> <li>• Begin process to establish an ecologically and societially acceptable level of risk for water availability and ecological health (Poff et al. 2009; Arthington et al. 2006).</li> <li>• Evaluate acceptable risk as a “balance between the perceived value of ecological goals, the economic costs involved, and the scientific uncertainties in functional relationships between ecological responses and flow alteration.” (Poff et al. 2009)</li> </ul>	<p>Chapter 6.1</p> <p>Appendix A</p> <p>Appendix J</p> <p>Appendix J</p> <p>Appendix J</p>
Implementation Options	<p>Conceptual plan for a basin-wide comprehensive water resource plan.</p> <p>Review of computer based tools for determining impacts of flow alteration.</p> <ul style="list-style-type: none"> <li>• Comprehensive water use plan – sustainable use of the Potomac River now and into the future, with opportunities for adaptive management.</li> <li>• Computer-based tools to predict impacts on flow and ecology from land and water use decisions</li> </ul>	<p>Chapter 6.3</p> <p>Appendix K</p> <p>Appendix K</p>

## 7.4. Lessons Learned

The MPRWA resulted in a number of lessons learned about the methods and approaches employed in the study. Insights arose from both the large river and stream/small river components and concern the steps involved in building the hydrologic foundation, model adequacy, projections of future conditions, and the stakeholder participation process. The Middle Potomac large river component applied the ESWM approach that has been used in several other watersheds including the Susquehanna River. It is driven by literature results and expert judgment primarily because of a lack of biological and/or hydrological data specific to the study area. The stream and small river component is one of the first-ever attempts to perform a large-scale quantitative analysis using the ELOHA methodology. This approach is more data driven.

A substantial effort is required to construct a hydrologic foundation that, despite inherent limitations and uncertainties, performs adequately in terms of both simulated daily flow time series and calculated flow metrics.

Model verification and flow metric testing are essential steps in building confidence in the simulated flow time series. Detailed comparison of simulated to observed results enhances confidence that the modeled time series and flow metrics are an acceptable representation of reality. In particular, flow metrics should behave as expected and relate to biota in an understandable way. Flow metrics should preferably represent portions of the hydrograph that can be influenced by management.

It is impossible to predict accurately what future conditions will look like. Evaluation of multiple future scenarios is useful for bounding the range of possible flow impacts. The range of conditions can then be utilized for planning and management purposes and adapted over time as additional information becomes available.

A modification of the stream classification approach described by Poff et al. (2010) proved best at delineating regionally distinct macroinvertebrate communities. The ELOHA framework recommends using similarities in baseline, or undisturbed, hydrologic regimes<sup>13</sup> to classify streams into types such as groundwater-fed streams, seasonally predictable snowmelt rivers, etc. These classes are expected to show distinct differences in their ecological characteristics and biological responses to flow alteration. In this study, stream classes were established based on similarities in the macroinvertebrate communities of reference stream sites, not in their hydrologies. Reference stream sites have good water quality, excellent habitat conditions, and are minimally disturbed by most anthropogenic factors. Their communities closely resemble the best possible natural state, and the streams—with few exceptions—have relatively undisturbed hydrologic regimes. The advantage of using reference sites rather than baseline hydrologies to classify streams is the impacts of multiple

---

<sup>13</sup> A second classification approach attempted in this study and described in Appendix D, section 4 (preliminary analysis) and Appendix G was the “attribute-based” approach. Stream classes are developed based on similarities in their current conditions (e.g., temperature, stream gradient, geology, and stream size) instead of similarities in their baseline hydrologic regimes. This approach does not require a modeling effort and assumes biological communities within a stream class will respond similarly when exposed to a hydrologic alteration (surface withdrawals were the hydrologic alteration tested in this study). The approach was not successful in the Middle Potomac study area (Table 7). The authors suspect this was because other, non-flow factors impact stream macroinvertebrates more than hydrologic alteration at many sampling sites.

anthropogenic factors are minimized, and the composition and species abundances of macroinvertebrate communities at the sites are primarily governed by natural factors. In a recursive partitioning analysis (a form of cluster analysis), reference communities separated most strongly on combinations of USEPA Level IV ecoregions. Therefore, stream classes in this study were established on aggregates of the ecoregions called bioregions.

The MPRWA study process engaged stakeholders at several points in the project. Stakeholder engagement was essential to conveying large quantities of technical information and receiving feedback.

### 7.5. Information Needs

The Middle Potomac River Watershed Assessment identified a number of information needs that could improve this assessment and/or benefit future work. These information needs fall into three categories: research and analysis, monitoring, and modeling.

#### 7.5.1. Research and Analysis

Recommendations made at the large river flow needs workshop (September 2010) focused primarily on monitoring needs. A few studies and analyses were suggested for mussel and fish populations, and ice scour impacts during low flows. A modified Instream Flow Incremental Method (IFIM) study was suggested to evaluate the relationship between flow and velocity, depth, and habitat below 1,000 MGD. The full set of suggestions can be found in Appendix A.

The large river flow needs study also identified a weakness in the existing literature. Specifically, there is a limited availability of information on flow requirements for aquatic species compared to velocity requirements. Flow is the total volume of water passing a fixed point over a given time period, and is commonly expressed as cubic feet per second (cfs) or million gallons per day (MGD). Flow is usually unaffected by stream morphology—speeding up when the stream narrows and slowing down when the stream widens. Velocity is the speed of the water, or the distance traveled by the water over time. It is strongly influenced by stream morphology. Translating flows into velocities for application at an ungauged location is difficult because detailed information about the depth, width, and slope at the new location are required.

Areas for further evaluation and analysis that were identified based on stakeholder feedback at the small river and stream flow needs workshop (November 2011) include evaluating (1) additional flow and macroinvertebrate metrics, (2) other taxa beyond benthic macroinvertebrates to develop flow-ecology relationships, (3) the results of the analyses using family versus genus level biological data, (4) flow-ecology relationships with additional stream classification factors, (5) the steep decline in stream health with less than 10 percent change in streamflow and the relationship of this to the model's margin of error (or, as was latter suspected, to a greater frequency of development-related, degraded water and habitat quality in the Piedmont), (6) the reliability of data at extreme ends of the FA-E curves, given the scarcity of data at the extremes, and (7) specific flow-ecology hypotheses. The full set of November 2011 stream and small river workshop participants' comments is available in Appendix J. Research needs identified by the project team during development of the stream flow ecology relationships include the need to enhance the understanding of (1) flow alteration in Coastal Plain watersheds and karst watersheds, (2) the confounding influences of non-flow factors

impacting ecosystems and biological communities, and (3) methods used to ameliorate the negative impacts of both low and high flow conditions (e.g. efficacy of best management practices). An overarching research need for both rivers and streams in the Middle Potomac watershed is a better understanding of aquatic and riparian community responses to naturally prolonged low flows associated with regional droughts.

### 7.5.2. Monitoring

Observed flow and biological data are essential to understanding and defining environmentally sustainable flows. Funding for sustained river monitoring is a critical priority. Specific data needs identified in both the small and large river studies are documented below.

The large river flow needs assessment identified a number of monitoring suggestions. Recommendations included (but were not limited to) long-term floodplain vegetation monitoring, continued/enhanced mussel and fish studies, and additional surveys related to cormorants, macroinvertebrates, and amphibians and reptiles. The complete, detailed list of recommendations is available in Appendix A.

Creating the hydrologic and biologic foundations upon which the FA-E relationships were built in the small streams assessment requires copious amounts of observed data. In using the existing data, a number of data needs were identified. Continued long-term operation of the Potomac basin stream gage network is the cornerstone of effective hydrologic model calibration and verification as well as flow metric verification efforts. Without these data, modeled flows cannot be tested. The locations of existing gages, however, are most often on medium and large sized waterways in non-karst regions. To fully understand hydrologic alteration and the associated ecological response throughout the Middle Potomac, additional observed streamflow data is needed on smaller streams, in karst watersheds, and in relatively undisturbed, or reference, watersheds.

Enhancements to existing biological datasets would enable a more diverse assessment of biological health. The Potomac is fortunate to have a basin-wide macroinvertebrate database; however, environmental flow analyses would also benefit from basin-wide datasets of other organisms such as fish, mussels, and vegetation because macroinvertebrates can only represent some elements of aquatic ecosystem health. For example, low magnitude flows do not appear to significantly impact macroinvertebrates whereas the flow-ecology literature suggests they might impact fish communities. Additional basin-wide, comparable datasets for fish could provide a more complete picture of the ecological impacts of flow alteration.

That said, two major hurdles impede the use of fish survey data for flow analyses. First, state and local monitoring programs typically use different methodologies to collect fish. Sampling protocols vary depending on management objectives. Constraints on time, personnel, and objectives can make it impractical to obtain standardized fish population estimates at all survey sites within a state (J. Mulligan, MDDNR, personal communication). Careful consideration of each program's methods is required to ensure the dataset assembled for flow analysis contains comparable data. Second, biological factors unrelated to flow or the stream environment significantly stress and alter fish populations in the Middle Potomac study area, including fish stocking programs, introductions of non-native species, and disease. A major analytical challenge will be determining the extent to which these biological factors, in addition to the known water quality and habitat problems, confound the impacts of flow alteration on fish populations.

### 7.5.3. Modeling

The hydrologic model utilized in this study has undergone substantial calibration and verification. These efforts are limited by the availability and nature of observed datasets, as previously noted. Due to limited observed data in small watersheds, karst watersheds, and reference watersheds, the ability to evaluate model behavior in these locations is somewhat compromised. As mentioned above, additional daily flow data in these areas could inform further model evaluation and enhancements.

Another information gap in the modeling effort relates to groundwater withdrawals. Although the largest withdrawals in the Potomac basin are taken from the surface water resources, there is a much larger number of groundwater withdrawals. The cumulative impacts of these groundwater withdrawals is likely significant in some areas. To this end, the hydrologic model either needs to be enhanced to be capable of effectively simulating these groundwater withdrawals or information needs to be obtained on the interaction of each groundwater withdrawal with surface water supplies for input into the existing hydrologic model.

Meteorological data drive the hydrologic cycle in the model. The broad spatial resolution of these meteorological data is a limitation to the simulation of smaller watersheds. Obtaining and importing high resolution meteorological data would likely improve estimation of flows, particularly in smaller watersheds where biological monitoring often occurs.

Implementation of management practices as a result of this study may be informed by more detailed modeling of the impacts of spatially explicit land use practices. For example, the HSPF model utilized in this study is not capable of spatially discriminating land uses within a land-river segment. However, the efficacy of management practices depends on their implementation location, among other factors. A modeling tool that can evaluate how site-specific management practices will affect stream ecology and stream health would be a useful addition to this analysis.

### 7.6. Applicability to Jurisdictions

The case studies and decision support tools described in Appendix K (DST\_CaseStudies.pdf) and in Kendy et al. (2012) demonstrate that states and river basins in the United States are successfully developing FA-E relationships and applying them to water resources planning and streamflow management. In lieu of the large hydrologic and biological datasets needed to develop FA-E curves, Richter et al. (2011) suggest implementing a presumptive flow standard based on the Sustainability Boundary Approach (Richter 2009). The presumptive standard restricts hydrologic alterations to within a percentage-based range of flows around natural or historic flow variability and protects vulnerable rivers from unlimited exploitation.

For the Middle Potomac study area, presumptive standards may best serve those watersheds that models cannot represent well at this time (Appendix D, section 2). For areas where models provide good representation, the FA-E relationships developed in the study (5.2, Appendix I) are generally applicable, with consideration for the confounding influences of other environmental factors and the innate differences in biometric response strength. All FA-E relationships show that macroinvertebrate family-level metrics respond as flow deviates from a baseline condition. Given the existing water quality and stream habitat conditions in the Middle Potomac region, a doubling in

flashiness (100 percent alteration) can be expected to result in roughly a 40 percent drop in acceptable biometric status and more than two thirds of stream communities—as measured by the multi-metric Chessie BIBI—in a degraded state (Figure 10). The precise change in the percent of degraded communities depends on whether concomitant changes in water quality and habitat conditions occur as flashiness increases. A 50 percent increase in the highest annual flow magnitudes could be expected to drop acceptable levels of biometrics by about 40 percent (Figure 6). Shortening high and low flow events to half (-50 percent) of their baseline durations can reduce acceptable biometric levels by about 35 percent (Figure 7a-b, 8). High flow events that occur twice as often (100 percent) can reduce acceptable biometric levels by roughly 20 percent (Figure 9). The successful development of FA-E relationships in the Middle Potomac study demonstrates that existing and planned changes in a watershed’s baseline flow regime can be related to alteration in streamflow and ultimately to responses in biometrics (Figure 12a-e).

There are potential applications of the study results for local and state jurisdictions. For example, a local water resources manager or planner might go through the following series of steps using FA-E relationships. At the local level, the results may be most applicable for land use planning and zoning.

- Identify the causes of ecological degradation in a watershed or jurisdiction of interest (e.g. flow alteration, water quality, stream habitat, etc.), and opportunities to mitigate those causes of degradation. This may be accomplished through various watershed assessment methodologies (e.g. Smith and Klimas 2013; USEPA 2000). If flow alteration is the primary cause of ecological degradation, proceed with the following steps. Otherwise, the flow alteration – ecological response relationships from this study would need to be modified before use.
- Determine how much ecological degradation is socially/politically acceptable within the area of interest. From a county perspective, a state perspective, and the Potomac River and Chesapeake Bay perspectives, how many degraded streams are allowable in the area of interest?
- Determine the watershed driver(s) of flow alteration in the area of interest (e.g. withdrawals, discharges, impoundments, land uses).
- For the watershed driver(s) of concern, determine how much of that driver will produce the maximum acceptable amount of degradation utilizing the modeled flow alteration – ecological response plots.
- Utilize regulatory, voluntary, or policy mechanisms to manage the watershed driver and prevent degradation past the acceptable threshold.

Similarly, state planning and regulatory entities may find the resulting FA-E relationships useful for state purposes, where withdrawal and discharge permitting typically occurs.

### 7.7. Next Steps

The Middle Potomac River Watershed Assessment establishes a scientific foundation for the larger process of identifying, managing and protecting environmental flows. Although information gaps and monitoring needs remain and should be addressed, social and political discussion and decision-making are important “next steps.” They might include:

## Middle Potomac River Watershed Assessment

- Use the study results to inform further development of the Potomac Basin Comprehensive Water Resources Plan and individual state plans. ICPRB is actively pursuing funding and stakeholder interest in order to pursue the development of this plan. USACE would provide supporting expertise in shared vision planning and stakeholder involvement. **ICPRB lead with USACE support.**
- Continue work begun in February 2012, when a technical workgroup was formed, under the auspices of the Maryland Department of Natural Resources (MDNR). Develop a hydroecological monitoring plan and list of priority research needs for the mainstem Potomac River. Other recommendations from the September 2010 Workshop for this workgroup, including developing more quantitative flow recommendations and re-evaluating the 300/100 mgd flow-by recommendation, have been deferred until such time as additional data have been gathered indicating adverse impacts on aquatic species due to flow alteration. **MDNR lead.**
- Consider how resource management agencies at local and state level in each jurisdiction can use the relationships developed in this assessment between land and water uses, flow alteration, and ecological status to inform watershed management decisions. This might include development of computer-based tools that use flow alteration-ecology relationships to predict the impact of proposed land and water use decisions. **TNC lead.**
- Use the quantitative flow alteration-ecology relationships to foster discussion and build consensus on acceptable levels of biological degradation resulting from changes in the flow regime. **TNC lead.**
- Development of a computer-based evaluation tool could facilitate understanding of the implications of various land and water use management decisions. **USACE** through the Interagency and International Support Technical Assistance Program.
- Perform additional research, monitoring, or analysis to fill information gaps and refine the findings from this study in an adaptive management context. **ICPRB and MDNR lead.**
- Use project data products to support other planning and management needs of basin jurisdictions. **TNC, ICPRB, and USACE.**

It is anticipated that these actions will be led by ICPRB, TNC, or basin jurisdictions. USACE may be a participating stakeholder in further activities, provide technical assistance, and will continue to advance the Chesapeake Bay restoration goals of Executive Order 13508 (The Chesapeake Bay Protection and Restoration Executive Order). No actions were identified for immediate USACE implementation.

## Supplemental Tables

### Supplemental Table A. Flow metrics.

Calculations were performed using the TNC Indicators of Hydrologic Alteration software program (IHA), the USGS Hydrologic Index Tool software program (HIT), the EPA DFLOW program, and with Excel 2007.

Metric Name	Description	Units
1-Day Maximum	The average of each year's highest daily mean flow (cfs) during the study period (1984-2005) divided by watershed area (mi <sup>2</sup> ). IHA.	cfs/mi <sup>2</sup>
3-Day Maximum	The average of each year's highest 3-day moving average of daily mean flow (cfs) during the study period (1984-2005) divided by watershed area (mi <sup>2</sup> ). IHA.	cfs/mi <sup>2</sup>
Annual Mean	The average of all the annual means of daily mean flows (cfs) during the study period (1984-2005) divided by watershed area (mi <sup>2</sup> ). The average of each year's mean daily flows is calculated, and then the means of each year are averaged. IHA.	cfs/mi <sup>2</sup>
Median	The median of all the daily mean flows (cfs) during the study period (1984-2005) divided by watershed area (mi <sup>2</sup> ). Excel.	cfs/mi <sup>2</sup>
Q85Seas	Calculate the 15th percentile flow (cfs) for each month (July – October) during the study period (1984-2005), and then average all the monthly values and divide by watershed area (mi <sup>2</sup> ). Adaptation of the Maryland method. Excel macro.	cfs/mi <sup>2</sup>
August Median	The median of the August median flow for each year in the study period (1984-2005) divided by watershed area (mi <sup>2</sup> ). IHA.	cfs/mi <sup>2</sup>
Base Flow Index	The median of each year's 7-day minimum flow (cfs) divided by the mean annual flow (cfs). IHA.	ratio (unitless)
1-Day Minimum	The average of each year's minimum daily mean flow (cfs) during the study period (1984-2005) divided by watershed area (mi <sup>2</sup> ). IHA.	cfs/mi <sup>2</sup>
3-Day Minimum	The average of each year's lowest 3-day moving average of daily flow (cfs) during the study period (1984-2005) divided by watershed area (mi <sup>2</sup> ). IHA.	cfs/mi <sup>2</sup>
7Q10	The lowest stream flow for seven consecutive days that would be expected to occur once in ten years. The 7-day moving average is calculated for the study period; the flow volume of the event that recurs every 10 years is the 7Q10 value. DFLOW.	cfs
High Pulse Duration	The median of the annual average number of consecutive days per year that daily flow is above the 90th percentile of the 1984-2005 period of record. IHA.	days/year

**Supplemental Table A. Continued.**

<b>Metric Name</b>	<b>Description</b>	<b>Units</b>
High Flow Index MH21	The average volume of high flow events (above a threshold equal to the median flow of the entire record) divided by the median daily flow for the entire record. HIT.	days
High Flow Duration DH17	The average duration of flow events with flows above the median flow for the entire period of record. HIT.	days
Flood Free Season	The length of the longest period common to all water years in the study period (1984-2005) where flows are at or below the high pulse threshold (usually less than the 90th percentile) in every year. IHA.	days
Low Pulse Duration	The median of the annual average number of consecutive days per year that daily flow is below the 10th percentile of the 1984-2005 period of record. IHA.	days/year
Extreme Low Flow Duration	Mean of extreme low flow event duration. An extreme low flow event is the occurrence of flow in the lowest 10th percentile of the low flows which are the lowest 10th percentile of all flows over the study period. IHA.	days/event
High Pulse Count	The median of the annual average of each year's number of times the daily mean flow is above the 90th percentile of all flows for the study period. IHA.	#/year
High Flow Frequency	Average of the number of events per year when the daily mean flow exceeds the 90th percentile of all flows in the study period. IHA.	#/year
Number of Reversals	The average number of times in a year that daily mean flow switches from rising to falling and vice versa. IHA.	#/year
Low Pulse Count	The median of the annual average of each year's number of times the daily mean flow is below the 10th percentile of all flows for the study period. IHA.	#/year
Extreme Low Flow Frequency	The frequency of extreme low flow events in a year, where daily flow is in the lowest 10th percentile of all the low flows (or below the 2.5th percentile of all flows in the 1984-2005 period of record). IHA.	#/year
Flashiness	(Richards-Baker Index) Sum of the absolute values of day-to-day changes in the daily mean flow divided by the sum of the daily mean flows. Excel.	ratio (unitless)
Rise Rate	The average of all positive differences in daily mean flow during "rising periods," or consecutive days for which change in daily flow is positive, in a year. IHA.	cfs/mi <sup>2</sup> /day
Fall Rate	The average of all negative differences in daily mean flow during "falling periods," or consecutive days for which change in daily flow is negative, in a year. IHA.	cfs/mi <sup>2</sup> /day

**Supplemental Table B. Macroinvertebrate family-level metrics.**

Calculations were performed using the Chesapeake Bay Program software program (Appendix 3). Metric type: C = composition; T = tolerance; R = richness; FG = feeding group; H = habit; index = multi-metric index. A metric can be more than one type. Family-level pollution tolerance values were originally developed by Hilsenhoff (1988) and refined by CBP. Values range from 0 to 10, with 0 being the most sensitive to pollution and 10 being the least (Appendix C).

Metric Name	Type	Description
ASPT Modified Index	T	Average of the family-level tolerance score of each family present in sample
Beck's Index	T	Index is $((3 \times \text{\#families with tolerance value of 0}) + (2 \times \text{\#families with tolerance value of 1}) + (1 \times \text{\#families with tolerance value of 2}))$
BIBI	index	Chesapeake Bay basin-wide Benthic Index of Biological Integrity; also called "Chessie BIBI;" average of scores of five bioregion-specific, family-level metrics
Gold Index	C	Index is 1 minus the proportional abundances (percents) of gastropods (snails), oligochaetes (segmented worms), and Diptera (true flies) individuals
Hilsenhoff Family Biotic Index	T	Also called FBI; average of the family-level tolerance score of each individual
# Ephemeroptera Families	R	Number of Ephemeroptera (mayflies) families present
# Sensitive Taxa	T	Number of families with family-level tolerance values less than or equal to 3
%Chironomidae	C	Percent of individuals that are Chironomids (non-biting midges)
%Clinger	H	Percent of individuals present that are adapted for clinging to hard surfaces
%Collector	FG	Percent of individuals that are collectors (filterers + gatherers) and not predatory
%Dominant3	T	Percent of individuals in the three most common families
%Ephemeroptera	C	Percent of individuals that are Ephemeroptera (mayflies)
%EPT	C	Percent of individuals that are Ephemeroptera (mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies)
%Filterer	FG	Percent of individuals that are adapted for filtering fine particles from the water column
%Gatherer	FG	Percent of individuals that are adapted for gathering a range of food particle sizes

**Supplemental Table B. Continued.**

<b>Metric Name</b>	<b>Type</b>	<b>Description</b>
%Scraper	FG	Percent of individuals that are adapted for scraping periphyton (algae, bacteria) from hard surfaces
%Swimmer	H	Percent of individuals that are adapted for swimming
%Tolerant	T	Percent of individuals with family-level tolerance values greater than or equal to 7
Shannon Wiener Index	R	Index is a measure of taxonomic diversity; it is the proportion of each family times the log of its proportion, summed for all families
Taxa Richness	R	The number of family-level taxa in the sample

## Literature Cited

- Apse, C., DePhilip, M., Zimmerman, J., and Smith, M. P. 2008. Developing instream flow criteria to support ecologically sustainable water resource planning and management. Final report to the Pennsylvania Instream Flow Technical Advisory Committee. The Nature Conservancy, Harrisburg, PA.
- Archfield, S.A., R.M. Vogel, P.A. Steeves, S.L. Brandt, P.K. Weiskel, and S.P. Garabedian. 2010. The Massachusetts Sustainable-Yield Estimator: A Decision-support Tool to Assess Water Availability at Ungaged Stream Locations in Massachusetts. Scientific Investigations Report. USGS.
- Astin, L. E. 2006. Data synthesis and bioindicator development for nontidal streams in the interstate Potomac River basin, USA. *Ecological Indicators*, 6: 664-685.
- Baker, D. B., R. P. Richards, T. T. Loftus, and J. W. Kramer. 2004. A new flashiness index: characteristics and applications to Midwestern rivers and streams. *Journal of the American Water Resources Association* 40(2):503-522.
- Bolger, R., R. Monsma, and R. Nelson. 2009. Sustainable Water Systems: Step One - Redefining the Nation's Infrastructure Challenge. Aspen Institute.
- Boulton, A. J. 2003. Parallels and contrasts in the effects of drought on stream macroinvertebrate assemblages. *Freshwater Biology* [Freshwat. Biol.]. Vol. 48 48:1173–1185.
- Brandes, D. G. J. Cavallo, and M. L. Nilson. 2005. Base flow trends in urbanizing watersheds of the Delaware River Basin. *Journal of the American Water Resources Association* 41(6):1377-1391.
- Brekke, L. D., J. E. Kiang, J. R. Olsen, R. S. Pulwarty, D. A. Raff, D. P. Turnipseed, R. S. Webb, and K. D. White. 2009. Climate Change and Water Resources Management - A Federal Perspective. Page 63. USGS, Reston, VA.
- Buchanan, C., K. Foreman, J. Johnson, and A. Griggs. 2011. Development of a Basin-Wide Benthic Index of Biotic Integrity for Non-Tidal Streams and Wadeable Rivers in the Chesapeake Bay Watershed: Final Report to the Chesapeake Bay Program Non-Tidal Water Quality Workgroup. ICPRB Report 11-1. Prepared for the US Environmental Protection Agency, Chesapeake Bay Program. <http://www.potomacriver.org/cms/publicationspdf/ICPRB11-01.pdf>.
- Bunn, S. E., and A. H. Arthington. 2002. Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environmental management* 30:492–507.

## Middle Potomac River Watershed Assessment

- Carlisle, D. M., and M. R. Meador. 2007. A biological assessment of streams in the eastern United States using a predictive model for macroinvertebrate assemblages. *Journal of the American Water Resources Association* 43(5): 1194-1207.
- Carlisle, D. M., D. M. Wolock, and M. R. Meador. 2010. Alteration of streamflow magnitudes and potential ecological consequences: a multiregional assessment. *Frontiers in Ecology and the Environment*. Doi:10.1890/100053.
- Chapra, S.C. 1997. *Surface water-quality modeling*. WCB McGraw-Hill, New York.
- Chesapeake Bay Program. 2010. Chesapeake Bay Program Environmental Modeling - Backgrounder. Annapolis, MD.  
<http://www.chesapeakebay.net/about/programs/modeling/53>.
- Cincotta, D. A., K. L. Dull, S. L. Markham, and R. D. Williams. 1986. Ichthyofaunal checklist and comparison of two North Branch Potomac River tributaries where alleged Ohio-Potomac River stream captures occurred. *Proceedings of the Pennsylvania Academy of Science* PPASAK 60(2):129-134.
- Cottingham P., Quinn G., Norris R., King A., Chessman B. and Marshall C. (2005). *Environmental Flows Monitoring and Assessment Framework*. Technical report. Cooperative Research Centre for Freshwater Ecology, Canberra.  
[http://freshwater.canberra.edu.au/publications.nsf/0/b217ed362dc90bca256fc7001cf693/\\$FILE/env%20flows%20monit%20assess%27t%20fr%27wrk.pdf](http://freshwater.canberra.edu.au/publications.nsf/0/b217ed362dc90bca256fc7001cf693/$FILE/env%20flows%20monit%20assess%27t%20fr%27wrk.pdf)
- Cruise, J. F., C. A. Laymon, and O. Z. Al-Hamdan. 2010. Impact of 20 years of land-cover change on the hydrology of streams in the southeastern United States. *Journal of the American Water Resources Association* 46(6):1159-1170.
- Cuffney, T. F., R. A. Brightbill, J. T. May, and I. R. Waite. 2010. Responses of benthic macroinvertebrates to environmental changes associated with urbanization in nine metropolitan areas. *Ecological Applications* 20(5):1384-1401.
- Cummins, J., C. Buchanan, C. Haywood, H. Moltz, A. Griggs, C. Jones, R. Kraus, N. P. Hitt, and R. Bumgardner. 2011. *Potomac Basin Large River Environmental Flow Needs*. ICPRB Report 10-3. Interstate Commission on the Potomac River Basin. 216 pages.
- De Waele, J., L. Plan, and P. Audra. 2009. Recent developments in surface and subsurface karst geomorphology: An introduction. *Geomorphology* 106:1-8.
- Dixon, M.D. 2003. Effects of flow pattern on riparian seedling recruitment on sandbars in the Wisconsin River, Wisconsin, USA. *Wetlands* 23:125-139.
- Donigian, A.S. 2002. Watershed model calibration and validation-the HSPF experience. In *Specialty Conference Proceedings: Water Environment Foundation*, 30. Phoenix, AZ.

## Middle Potomac River Watershed Assessment

- Dunne, T., and L. Leopold. 1978. *Water in environmental planning*. W.H. Freeman and Company, New York.
- Everson, D. A., and D. H. Boucher. 1998. Tree species-richness and topographic complexity along the riparian edge of the Potomac River. *Forest Ecology and Management* 109:305-314.
- Falcone, J. A., D. M. Carlisle, and L. C. Weber. 2012. Quantifying human disturbance in watersheds: variable selection and performance of a GIS-based disturbance index for predicting the biological condition of perennial streams. *Ecological Indicators* 10:264-273.
- Fausch, K. D., Y. Taniguchi, S. Nakano, G. D. Grossman, and C. R. Townsend. 2001. Flood disturbance regimes influence rainbow trout invasion success among five holarctic regions. *Ecological Applications* 11: 1438-1455.
- Feminella, J. W. 2000. Correspondence between stream macroinvertebrate assemblages and 4 ecoregions of the southeastern USA. *Journal North American Benthological Society* 19(3):442-461.
- Foreman, K., C. Buchanan, and A. Nagel. 2008. Development of ecosystem health indexes for non-tidal wadeable streams and rivers in the Chesapeake Bay basin. Unpublished report to the Chesapeake Bay Program Non-Tidal Water Quality Workgroup. December 5, 2008.
- Freeman, M. C., and P. A. Marcinek. 2006. Fish assemblage responses to water withdrawals and water supply reservoirs in Piedmont streams. *Environmental Management* 38:435-450.
- Gassman, P.W., M.R. Reyes, C.H. Green, and J.G. Arnold. 2007. *The soil and water assessment tool: Historical development, applications, and future research directions*. Ames, Iowa: Center for Agricultural and Rural Development, Iowa State University.
- Gordon, N. D., T. A. McMahon, and B. L. Finlayson. 1992. *Stream Hydrology: an Introduction for Ecologists*. John Wiley & Sons, 542 pp.
- Hakala, J. P. and K. J. Hartman. 2004. Drought effects on stream morphology and brook trout (*Salvelinus fontinalis*) populations in forested headwater streams. *Hydrobiologia* 515:203-213.
- Hamon, W.R. 1961. Estimating potential evapotranspiration. *Proceedings of the American Society of Civil Engineers* 87:107-120.
- Hawkins, C. P., R. H. Norris, J. Gerritsen, R. M. Hughes, S. K. Jackson, R. K. Johnson, and R. J. Stevenson. 2000. Evaluation of the use of landscape classifications for the prediction of freshwater biota: synthesis and recommendations. *Journal North American Benthological Society* 19(3):541-556.
- Henriksen, J. A., Heasley, J., Kennen, J.G., and Niewsand, S. 2006. Users' manual for the hydroecological integrity assessment process software (including the New Jersey Assessment Tools): U.S. Geological Survey, Biological Resources Discipline, Open File Report 2006-1093, 71 p. <http://www.fort.usgs.gov/Products/Software/NATHAT/>

## Middle Potomac River Watershed Assessment

- Hilsenhoff, W. L. (1988). Rapid field assessment of organic pollution with a family-level biotic index. *J. North American Benthological Society*: 7(1):65-68.
- Hitt, N. P., and P. L. Angermeier. 2008. Evidence for fish dispersal from spatial analysis of stream network topology. *Journal of the North American Benthological Society* 27: 304-320.
- Hocutt, C. H. 1979. Drainage evolution and fish dispersal in the central Appalachians: summary. *Geological Society of America Bulletin* 90:129.
- Holomuzki, J. R., and B. J. F. Biggs. 2000. Taxon-Specific Responses to High-Flow Disturbance in Streams: Implications for Population Persistence. *Journal of the North American Benthological Society* 19:670. doi: 10.2307/1468125.
- Humphries, P., and D. S. Baldwin. 2003. Drought and aquatic ecosystems: an introduction. *Freshwater Biology* 48:1141–1146. doi: 10.1046/j.1365-2427.2003.01092.x.
- Hupp, C. R., and W. R. Osterkamp. 1985. Bottomland vegetation distribution along Passage Creek, Virginia, in relation to fluvial landforms. *Ecology* 66(3):670-681.
- Hynes, H. B. N. 1970. *The ecology of running waters*. University of Toronto Press. 555 pp.
- Jenkins, R. E., and N. M. Burkhead. 1994. *Freshwater Fishes of Virginia*. American Fisheries Society, Bethesda, Maryland.
- Kendy, E., C. Apse, and K. Blann. 2012. *A practical guide to environmental flows for policy and planning with nine case studies in the United States*. The Nature Conservancy.
- Kennan, J. G. 1999. Relation of macroinvertebrate community impairment to catchment characteristics in New Jersey streams. *Journal of the American Water Resources Association* 35(4):939-955.
- Kennan, J. G., K. Riva-Murray, and K. M. Beaulieu. 2009. Determining hydrologic factors that influence stream macroinvertebrate assemblages in the northeastern US. *Ecology* DOI: 10.1002/eco.
- Kenny, Joan F., Nancy L. Barber, Susan S. Hutson, Kristin S. Linsey, John K. Lovelace, and Molly A. Maupin. 2009. *Estimated Use of Water in the United States in 2005*. Scientific Investigations Report. USGS. <http://pubs.usgs.gov/circ/1344/>.
- Kiang, J., and E. Hagen. 2003. 2002 drought operations and lessons learned: Washington metropolitan area. Interstate Commission on the Potomac River Basin.
- Konrad, C. 2011. Principal technical decision points for developing regional environmental flow standards. Unpublished draft, March 1, 2011.
- Kudlas, S. 2009. Models, Uncertainty, and Cumulative Impacts on Water Resources: The Virginia DEQ Approach to Water Resource Modeling presented at the Southeast States Water

## Middle Potomac River Watershed Assessment

- Resources Management and Planning Forum, Richmond, VA.  
[http://sogweb.sog.unc.edu/Water/images/7/72/Vadeq\\_approach\\_v02-Kudlas.pdf](http://sogweb.sog.unc.edu/Water/images/7/72/Vadeq_approach_v02-Kudlas.pdf).
- Lake, P. S. 2003. Ecological effects of perturbation by drought in flowing waters. *Freshwater Biology* 48:1161–1172. doi: 10.1046/j.1365-2427.2003.01086.x.
- Lawrence, R., and A. Wright. 2001. Rule-based classification system using Classification and Regression Tree (CART) Analysis. *Photogrammetric Engineering and Remote Sensing* 67:1137-1142.
- Legrand, H., and V. Stringfield. 1973. Karst hydrology - A review. *Journal of Hydrology* 20:97-120.
- Linker, L. C., G. W. Shenk, R. L. Dennis, and J. S. Sweeney. 1999. Cross-media models of the Chesapeake Bay watershed and airshed. Annapolis, MD: Chesapeake Bay Program Office.  
<http://archive.chesapeakebay.net/pubs/948.pdf>.
- Lumb, A.M., R.B. McCammon, and J.L. Kittle. 1994. Users manual for an Expert System (HSPEXP) for calibration of the Hydrological Simulation Program: Fortran. Water Resources Investigations Report. Reston, VA: USGS.
- Lytle, D., and N. L. Poff. 2004. Adaptation to natural flow regimes. *Trends in Ecology & Evolution* 19:94–100. doi: 10.1016/j.tree.2003.10.002.
- Maryland Department of Natural Resources (MD DNR). 1981. Potomac River Environmental Flow-by Study. Maryland Department of Natural Resources. Baltimore, MD
- Merritt, R. W., K. W. Cummins, and M. B. Berg (eds). 2008. *An Introduction to the Aquatic Insects of North America*. Fourth Edition. Kendal/Hunt Publishing Co. 1214 pgs.
- Moyer, D., and M. Bennett. 2007. Development of relations of stream stage to channel geometry and discharge for stream segments simulated with Hydrologic Simulation Program - Fortran (HSPF), Chesapeake Bay Watershed and Adjacent Parts of Virginia, Maryland, and Delaware. USGS Scientific Investigations Report, USGS. Retrieved from  
<http://pubs.usgs.gov/sir/2007/5135/pdf/SIR2007-5135.pdf>.
- Nash, J.E., and J.V. Sutcliffe. 1970. River flow forecasting through conceptual models, Part 1: A discussion of principles. *Journal of Hydrology* 10:282-290.
- Novotny, V., and H. Olem. 1994. *Water Quality: Prevention, Identification, and Management of Diffuse Pollution*. Van Nostrand Reinhold, New York.
- Olden, J.D., and N.L. Poff. 2003. Redundancy and the choice of hydrologic indices for characterizing streamflow regimes. *River Research and Applications* 19:101-121. doi: 10.1002/rra.700.

- Olivero, A. P., and M. G. Anderson. 2008. Northeast aquatic habitat classification system. Report prepared by The Nature Conservancy Eastern Regional Office in collaboration with the Northeast Association of Fish and Wildlife Agencies.  
[http://www.fws.gov/filedownloads/ftp\\_gis/R5/GOMP/Jed/Prints/6/3%20NE%20Habitat%20Classification%20-%20Aquatic.pdf](http://www.fws.gov/filedownloads/ftp_gis/R5/GOMP/Jed/Prints/6/3%20NE%20Habitat%20Classification%20-%20Aquatic.pdf)
- Ortmann, A. E. 1913. The Alleghenian divide, and its influence upon the freshwater fauna. *Proceedings of the American Philosophical Society* 52: 287-381.
- PADEP (Pennsylvania Department of Environmental Protection). 2009. Commonwealth of Pennsylvania State Water Plan Principles. Page 82. Retrieved from <http://www.pawaterplan.dep.state.pa.us/docs/Publications/3010-BK-DEP4222.pdf>.
- Poff N.L. and Zimmerman J.K.H. 2010. Ecological responses to altered flow regimes: a literature review to inform environmental flows science and management. *Freshwater Biology* 55:194-20.
- Poff, N. L., and J. V. Ward. 1989. Implications of Streamflow Variability and Predictability for Lotic Community Structure: A Regional Analysis of Streamflow Patterns. *Canadian Journal of Fisheries and Aquatic Sciences* 46:1805–1818. doi: 10.1139/f89-228.
- Poff, N. L., B. D. Richter, A. H. Arthington, S. E. Bunn, R. J. Naiman, E. Kendy, M. Acreman, C. Apse, B. P. Bledsoe, M. C. Freeman, J. Henriksen, R. B. Jacobson, J. G. Kennen, D. M. Merritt, J. H. O’Keeffe, J. D. Olden, K. Rogers, R. E. Tharme, and A. Warner. 2010. The ecological limits of hydrologic alteration (ELOHA): a new framework for developing regional environmental flow standards. *Freshwater Biology* 55(1):147–170.
- Poff, N. L., J. D. Olden, N. K. M. Vieira, D. S. Finn, M. P. Simmons, and B. C. Kondratieff. 2006. Functional trait niches of North American lotic insects: traits-based ecological applications in light of phylogenetic relationships. *Journal of the North American Benthological Society* 25:730-755.
- Reeves, H.W., D.A. Hamilton, P.W. Seelbach, and A.J. Asher. In review. Ground-water Component of the Michigan Water Withdrawal Screening Tool. Scientific Investigations Report. USGS.
- Richardson, S. M., J. Hanson, and A. Locke. 2002. Effects of impoundment and water-level fluctuations on macrophyte and macroinvertebrate communities of a dammed tidal river. *Aquatic ecology [Aquat. Ecol.]*. Vol. 36:493–510.
- Richter BD. 2009. Re-thinking environmental flows: from allocations and reserves to sustainability boundaries. *Rivers Research and Applications* 25: 1–12.
- Richter, B.D., M. Davis, C. Apse, and C.P. Konrad. 2011. A presumptive standard for environmental flow protections. *River Research and Applications*. Doi: 10.1002/rra.1511.

## Middle Potomac River Watershed Assessment

- Richter, B.D., R. Mathews, D.L. Harrison, and R. Wigington, 2003. Ecologically sustainable water management: Managing river flows for ecological integrity. *Ecological Applications* 13:206-224.
- Robinson, C. T., S. Aebischer, and U. Uehlinger. 2004. Immediate and habitat-specific responses of macroinvertebrates to sequential, experimental floods. *Journal of the North American Benthological Society* 23:853–867. doi: 10.1899/0887-3593(2004)023<0853:IAHROM>2.0.CO;2.
- Rode, M., and U., Suhr. 2006. Uncertainties in selected surface water quality data. *Hydrology and Earth System Sciences* 3:2991-3021.
- Rossman, L.A. 1990. DFLOW User's Manual. Washington, DC: EPA.
- Rutledge, A. 1998. Computer programs for describing the recession of ground-water discharge and for estimating mean ground-water recharge and discharge from streamflow data - update. Page 43. Water Resources Investigations Report, USGS.
- SAS Institute Inc. 2009. SAS OnlineDoc®, version 9.2. SAS Institute Inc., Cary, NC.
- Schultz, C.L., and J. Palmer. 2008. Seasonal Steady-State Ground Water/Stream Flow Model of the Upper Monocacy River Basin. Page 46. Interstate Commission on the Potomac River Basin, Rockville, MD. Retrieved from [http://www.potomacriver.org/cms/drinkingwaterdocs/NFWF\\_GWProject/NFWF\\_UpperMonoc\\_GWModelDocumentation\\_Sep-30-2008.pdf](http://www.potomacriver.org/cms/drinkingwaterdocs/NFWF_GWProject/NFWF_UpperMonoc_GWModelDocumentation_Sep-30-2008.pdf).
- Schultz, C.L., D. Tipton, and J. Palmer. 2004. Annual and Seasonal Water Budgets for the Monocacy/Catoctin Drainage Area. Page 85. ICPRB, Rockville, MD. Retrieved from <http://www.potomacriver.org/cms/publicationspdf/ICPRB04-4.pdf>.
- Schultz, C.L., S.A. Ahmed, R. Mandel, and H. Moltz. In review. Implementation of a nonlinear groundwater storage-discharge algorithm in HSPF model of the upper Potomac River basin.
- Schwartz, F. J. 1956. The distribution and probable postglacial dispersal of the percid fish, *Etheostoma b. blennioides*, in the Potomac River. *Copeia* 1965(3): 285-290.
- Semmens, D., M., Hernandez, D.C. Goodrich, and W.G. Kepner. 2006. A retrospective analysis of model uncertainty for forecasting hydrologic change. Proceedings of iEMSs 3rd Biennial Meeting: Summit on Environmental Modeling and Software, International Environmental Modeling and Software Society, Burlington, VT.
- Sepkoski, J. J. and M. A. Rex. 1974. Distribution of Freshwater Mussels: Coastal Rivers as Biogeographic Islands. *Systematic Zoology* 23: 165-188.
- Sheldon, A. L. 1988. Conservation of stream fishes: patterns of diversity, rarity, and risk. *Conservation Biology* 2: 149-156.

## Middle Potomac River Watershed Assessment

- Smith, R.D., and C.V. Klimas. 2013. A multi-scale approach to assess and restore ecosystems in a watershed context. U.S. Army Corps of Engineers, Engineer Research and Development Center. ERDC/EL TR-13-18.
- SRBC (Susquehanna River Basin Commission). 2007. Pennsylvania Agricultural Consumptive Water Use. Page 2. Information Sheet, SRBC. Retrieved from [http://www.srbc.net/pubinfo/docs/Agricultural%20Water%20Use%20%281\\_07%29.PDF](http://www.srbc.net/pubinfo/docs/Agricultural%20Water%20Use%20%281_07%29.PDF).
- SRBC (Susquehanna River Basin Commission). 2008. Consumptive Use Mitigation Plan. Page 35. Susquehanna River Basin Commission, Harrisburg, PA. Retrieved from <http://www.srbc.net/programs/docs/CUMP.pdf>.
- Starnes, W. C. 2002. Current diversity, historical analysis, and biotic integrity of fishes in the lower Potomac basin in the vicinity of Plummerville Island, Maryland - Contribution to the natural history of Plummerville Island, Maryland XXVII. Proceedings of the Biological Society of Washington 115: 273-320.
- Steiner, R. C., E. R. Hagen, and J. Duncanson. 2000. Water Supply Demands and Resource Analysis in the Potomac River Basin. Interstate Commission on the Potomac River Basin.
- Steiner, R.C. 1999. Cooperative management of Washington area 1999 drought operations. Interstate Commission on the Potomac River Basin.
- Strahler, A.N. 1957. Quantitative analysis of watershed geomorphology. American Geophysical Union Transactions. 38:913-920.
- Stuckey, M. H. 2006. Low-flow, base-flow, and mean-flow regression equations for Pennsylvania Streams. U.S. Geological Survey, Scientific Investigations Report 2006-5130.
- Taylor, R. W. 1985. Comments on the distribution of freshwater mussels (Unionacea) of the Potomac River headwaters in West Virginia. Nautilus 99(2-3): 84-87.
- The Nature Conservancy. 2007. Indicators of Hydrologic Alteration, Version 7 User's Manual. <http://www.nature.org/initiatives/freshwater/conservationtools/>.
- Thompson, H. D. 1939. Drainage evolution in the southern Appalachians. Bulletin of the Geological Society of America 50(8): 1323-1356.
- Thorp, J. H., and A. P. Covich (eds). 2001. Ecology and Classification of North American Freshwater Invertebrates. Second Edition. Elsevier Science. 1056 pgs.
- U.S. Army Corps of Engineers. 2004. Middle Potomac Watershed, Maryland, Pennsylvania, Virginia, West Virginia, District of Columbia - Section 905(b) (WRDA 86) Analysis. Page 43. Baltimore, MD.
- U.S. Army Corps of Engineers. 2009a. Ohio River Basin Comprehensive Reconnaissance Report. Page 232. U.S. Army Corps of Engineers.

## Middle Potomac River Watershed Assessment

- U.S. Army Corps of Engineers. 2009b. Middle Potomac River Watershed Assessment, Potomac River Sustainable Flow and Water Resources Analysis: Project Management Plan. U.S. Army Corps of Engineers, Baltimore, MD.
- U.S. Army Corps of Engineers. 2010. Building Strong Collaborative Relationships for a Sustainable Water Resources Future. National Report: Responding to National Water Resources Challenges. Page 104. U.S. Army Corps of Engineers, Washington, DC. Retrieved November 1, 2010, from <http://www.building-collaboration-for-water.org/>.
- US EPA (United States Environmental Protection Agency). 2000. Stressor Identification Guidance Document. EPA/822/B-00/025. <http://www.epa.gov/caddis/index.html>.
- USEPA (U.S. Environmental Protection Agency). 2005. National Hydrography Dataset Plus – NHDPlus. Produced by Horizon Systems.
- USEPA (U.S. Environmental Protection Agency). 2007. BASINS Technical Note 1: Creating Hydraulic Function Tables (FTABLES) for Reservoirs in BASINS. U.S. Environmental Protection Agency. Retrieved from <http://www.epa.gov/waterscience/basins/docs/tecnote1.pdf>.
- USEPA (U.S. Environmental Protection Agency). 2009. Chesapeake Bay Phase 5 Community Watershed Model. U.S. Environmental Protection Agency, Chesapeake Bay Program Office, Annapolis, MD.
- USEPA (U.S. Environmental Protection Agency). 2009. Technical Guidance on Implementing the Stormwater Runoff Requirements for Federal Projects under Section 438 of the Energy Independence and Security Act. Page 61. EPA, Washington, CC. Retrieved from [www.epa.gov/owow/nps/lid/section438](http://www.epa.gov/owow/nps/lid/section438).
- USEPA (U.S. Environmental Protection Agency). 2010. Chesapeake Bay Phase 5.3 Community Watershed Model. EPA 903S10002 - CBP/TRS-303-10. U.S. Environmental Protection Agency, Chesapeake Bay Program Office, Annapolis MD. December 2010.
- Vann, D. T., R. Mandel, J. M. Miller, E. Hagen, A. Buda, and D. Cordalis. 2004. The District of Columbia Source Water Assessment. ICPRB, Rockville, MD.
- Vannote, R. L., G. W. Minshall, K. W. Cummins, J. R. Sedell, and C. E. Cushing. 1980. The River Continuum Concept. *Canadian Journal of Fisheries and Aquatic Sciences* 37:130-137.
- Venables, W.N., and D.N. Smith. 2011. An introduction to R: Notes on R: A programming environment for data analysis and graphics. R Development Core Team. .
- Waldron, M. C., and S. A. Archfield. 2006. Factors Affecting Firm Yield and the Estimation of Firm Yield for Selected Streamflow-dominated Drinking-water-supply Reservoirs in Massachusetts, US Scientific Investigations Report. USGS. <http://pubs.usgs.gov/sir/2006/5044/pdf/SIR2006-5044.pdf>. Ward, J. V. 1991. Aquatic Insect Ecology. Part I, Biology and Habitat. Wiley. 456 pgs.

## Middle Potomac River Watershed Assessment

- Ward, J. V. 1992. Aquatic insect ecology: 1. Biology and habitat. John Wiley & Sons, New York, 438p.
- Watters, G. T. and S. H. O'dee. 1998. Metamorphosis of Freshwater Mussel Glochidia (Bivalvia: Unionidae) on Amphibians and Exotic Fishes. *American Midland Naturalist* 139: 49-57.
- Weidner, E. 2009. Potomac Watershed Priority Lands Strategy: Conserving Lands to Benefit Drinking Water Quality. Duke University. Retrieved from [http://www.potomacdwsp.org/Meetings/Jul22-2009/Weidner\\_Report.pdf](http://www.potomacdwsp.org/Meetings/Jul22-2009/Weidner_Report.pdf).
- Weiskel, P.K., S.L. Brandt, L.A. DeSimone, L.J. Ostiguy, and S.A. Archfield. 2010. Indicators of Streamflow Alteration, Habitat Fragmentation, Impervious Cover, and Water Quality for Massachusetts Stream Basins. Scientific Investigations Report. USGS. <http://pubs.usgs.gov/sir/2009/5272/>.
- Wetzel, R. G. 2001. Limnology. Lake and River Ecosystems. Third Edition. Academic Press. 1006 pgs.
- Williamson, S., C. Horin, M. Ruth, R. F. Weston, K. Ross, and D. Irani. 2008. Climate Change Impacts on Maryland and the Cost of Inaction: A Review and Assessment by the Center for Integrated Environmental Research (CIER) at the University of Maryland. Page 24. Maryland Commission on Climate Change. Retrieved from <http://www.cier.umd.edu/climateadaptation/Chapter3.pdf>.
- Wood, A. J., J. M. Olmernik, and D. D. Brown. 1999. Level III and IV ecoregions of Delaware, Maryland, Pennsylvania, Virginia and West Virginia. [http://www.epa.gov/wed/pages/ecoregions/reg3\\_eco.htm](http://www.epa.gov/wed/pages/ecoregions/reg3_eco.htm).

## Glossary

**Baseline Scenario** – The set of model conditions applied to catchments that defined landscape, withdrawal, and discharge conditions that would result in an unaltered hydrologic regime.

**Bioregion/ecoregion/physiographic province** – Classification systems that defines geographic regions with similar attributes such as geology, soils, elevation, slope, and climate, and is useful for classifying ecological data for study.

**Classification and Regression Tree (CART) and Recursive Partitioning (RPART)** – Statistical methods that delineate groups and identify numerical breakpoints among independent variables and partition variance in the response variable.

**Hydrologic Simulation Program-FORTRAN (HSPF)** – Chesapeake Bay Program’s version of the HSPF modeling system adapted by the Chesapeake Bay Program and utilized in this project in combination with the WOOOMM to simulate streamflows at ungaged locations.

**Consumptive use** – The amount of water withdrawn that is not subsequently returned to the source.

**Current scenario** – The set of model conditions applied to catchments that defined landscape, withdrawal, and discharge conditions that result in the existing hydrologic regime.

**DFLOW** – Software developed by the EPA to estimate low flow metrics from daily input time series.

**Ecological Limits of Hydrologic Alteration (ELOHA)** – Methodology developed by The Nature Conservancy, ELOHA is a “scientifically robust and flexible framework for assessing and managing environmental flows across large regions, when lack of time and resources preclude evaluating individual rivers.”

**Environmental flows** – “Flows that sustain healthy river ecosystems and the goods and services that humans derive from them.”

**Flow Alteration-Ecology Response (FA-E)** – Measureable shifts of biological indicator communities in response to the stresses caused by altered flow regimes.

**Flow metric** – A statistical measure of a system’s hydrologic characteristic, based on an input flow time series.

**Geographic Information Systems (GIS)** – A GIS “integrates hardware, software, and data for capturing, managing, analyzing, and displaying all forms of geographically referenced information.” (ESRI)

**Hydrologic Index Tool (HIT)** – Software developed by USGS to calculate 171 ecologically relevant flow metrics from input daily flow time series.

**Hydrologic alteration** – The amount of change in a flow metric between two conditions.

**Indicators of Hydrologic Alteration (IHA)** – Software developed by The Nature Conservancy to “understand hydrologic changes in ecologically-relevant terms.” The software computes 67 flow metrics from input daily flow time series.

**Karst** – landscape formed by the dissolution of soluble carbonate bedrock such as limestone, dolomite, and gypsum. It has naturally elevated conductivity and is characterized by many springs, caves, sinkholes, and a network of aquifers and underground channels.

**Macroinvertebrates** – Animals without spines that are visible to the naked eye. Commonly includes insects, arthropods, and crustaceans.

**Model scenarios** – A run of a hydrologic model with a specific set of user-defined input data. In the Middle Potomac River Watershed Assessment, there are seven model scenarios (baseline, current, domestic and public supply 1 and 2, power, climate change, and hot and dry).

**Nash-Sutcliffe Efficiency** – A test capable of measuring the predictive power of a hydrologic model.

**Reference** – A series of biological sampling events that met an *a-priori* defined set of high-quality physical habitat and water quality conditions, which are expected to support high-quality biological communities.

**River Continuum Concept** – A general classification system for flowing waters and their communities from headwaters to large rivers. The concept is based on the idea that biological communities reach predictable dynamic equilibriums in response to physical river attributes of depth, width, velocity, and longitudinal shifts in productivity and respiration.

**River segment** – A uniquely modeled watershed area in the CBP HSPF model.

**Watershed Data Management (WMD)** – A binary input file used to store input time series for the HSPF model.

**Online Object Oriented Meta-Model (WOOOMM)** – a hydrologic routing module developed by Virginia Department of Environmental Quality and utilized in this project to route the CBP HSPF model to simulate streamflows at ungaged locations.

## Abbreviations and Acronyms

AG	agricultural sector
AWUD	Aggregate Water Use Data system
CART	Category and Regression Tree Analysis
CBP	Chesapeake Bay Program
cfs	cubic feet per second
cfs/mi <sup>2</sup>	cubic feet per second per square mile of watershed (water yield)
Chessie BIBI	Chesapeake Bay basin-wide Benthic Index of Biological Integrity
CO-OP	ICPRB section for cooperative water supply operations on the Potomac
CU	consumptive use
CUMP	consumptive use mitigation plan
DC	District of Columbia
DEM	digital elevation model
DP	domestic and public supply sector
DP1	domestic and public supply model scenario 1
DP2	domestic and public supply model scenario 2
DP3	domestic and public supply model scenario 3
ELOHA	Ecological Limits of Hydrologic Alteration
EPRI	Electric Power Research Institute
ESWM	Ecologically Sustainable Water Management
EVAP	evapotranspiration
FA-E	flow alteration-ecology relationship
FIPS	Federal Information Processing Standards
FRIS	Farm and Ranchland Information Survey
FTABLE	hydraulic function table
GIS	geographic information system
GMU	George Mason University
GW	groundwater
HD	hot and dry model scenario
HIT	hydrologic index tool
HSPF	Hydrological Simulation Program-FORTRAN
HUC	hydrologic unit code
ICPRB	Interstate Commission on the Potomac River Basin
IHA	indicators of hydrologic alteration
IN	industrial sector
IPCC	Intergovernmental Panel on Climate Change
IR	irrigation sector
IWRM	integrated water resources management
LV	livestock sector
MD	Maryland
MDDNR	Maryland Department of Natural Resources
MDE	Maryland Department of the Environment
MGD	million gallons per day
MG/year	million gallons per year
MI	mining sector

## Middle Potomac River Watershed Assessment

MPRWA	Middle Potomac River Watershed Assessment
MWD	multi-watershed delineation
NASS	National Agricultural Statistics Service
NEAFWA	Northeast Association of Fish and Wildlife Agencies
NEAHCS	Northeast Aquatic Habitat Classification System
NED	National Elevation Dataset
NHD	National Hydrography Dataset
NID	National Inventory of Dams
NOAA	National Oceanic and Atmospheric Administration
NPDES	National Pollution Discharge Elimination System
NPS	National Park Service
NRCS	Natural Resources Conservation Service
NSE	Nash-Sutcliffe Efficiency
PA	Pennsylvania
PADEP	Pennsylvania Department of Environmental Protection
PMP	Project Management Plan
PO	power sector
RESAC	Remote Sensing Application Center
RPART	Recursive Partitioning and Regression Tree
SRBC	Susquehanna River Basin Commission
SRES	IPCC special report on emission scenarios
SW	surface water
TAG	Technical Advisory Group
TMDL	total maximum daily load
TNC	The Nature Conservancy
USACE	United States Army Corps of Engineers
USDA	United States Department of Agriculture
USEIA	United States Energy Information Administration
USEPA	United States Environmental Protection Agency (also EPA)
USGS	United States Geological Society
VA	Virginia
VADEQ	Virginia Department of Environmental Quality
WAD	Washington Aqueduct
WD	withdrawal
WDM	Watershed Data Management format
WMA	Washington metropolitan area
WOOOMM	Online Object Oriented Meta-Model
WRE	water resources element
WSM	watershed model
WSSC	Washington Suburban Sanitary Commission
WV	West Virginia
WVDEP	West Virginia Department of Environmental Protection

## Team Members and Contributors

**ICPRB Team:** Carlton Haywood, Claire Buchanan, Heidi Moltz, Jim Palmer, Adam Griggs, Olivia Devereux, Karin Bencala

**ICPRB Contributors:** Ross Mandel, Jan Ducnuigeen, Cherie Shultz, Curtis Dalpra, Jacqueline Johnson, Jim Cummins

**TNC Team:** Stephanie Flack, Julie Zimmerman, Katherine Boomer

**USACE Team:** Claire O'Neill, Andrew Roach

## Acknowledgements

Rob Burgholzer, Virginia Department of Environmental Quality, made the WOOOMM module available to the project team and spent much time working with us on model inputs, model runs, and on testing model outputs. That assistance is much appreciated.

Tara Moberg and Michele DePhilip of Pennsylvania Chapter of The Nature Conservancy provided valuable advice on the Indicators of Hydrologic Alteration specifically and generally about the ESWM methodology, based on their experience doing similar work for the Susquehanna River.

This project benefited from the discussion of ideas, of early versions of products, and of draft conclusions, with participants at the two workshops and with members of the Technical Advisory Group. Those individuals are identified in the workshop summaries and in the Stakeholder involvement summary (Appendices A and J). Special thanks are extended to Eloise Kendy, Colin Apse, Doug Sampson, Julie Zimmerman, John Mullican, Matthew Ashton, Steve Schreiner, and Fairfax Water for their reviews of the draft report and valuable suggestions.

The basic tool used for flow modeling was the Chesapeake Bay Watershed Model. Gary Shenk, US EPA, answered many questions about running that model.

The United State Geological Survey and its partners collected the surface flow data at gaging stations in the Potomac and Susquehanna watersheds and disseminated the data to the public, state and local governments, public and private utilities, and other federal agencies involved with managing water resources.

Monitoring programs of federal, state, interstate, and local agencies and academic institutions in Delaware, Maryland, New York, Pennsylvania, Virginia, and West Virginia collected and processed the stream data made available to this project, and the Chesapeake Bay Program (CBP) incorporated the data into a uniform database and calculated the suite of biometrics used in the project.

# Appendix A – Potomac Large River Environmental Flow Needs Report

---

## Contents

This appendix contains the *Potomac Large River Environmental Flow Needs* report, which identifies the hydrologic needs of flow-dependent species and communities in four segments of the mainstem Potomac River and two selected large tributaries. The report was developed by a research team from the Interstate Commission on the Potomac River Basin (ICPRB), the Nature Conservancy (TNC), Leetown Science Center Aquatic Ecology Branch of the U.S. Geological Survey (USGS), and the Potomac Environmental Research and Education Center of George Mason University (GMU). It includes a comprehensive literature review, development of flow hypotheses, assessment of large river environmental flow needs, statistics proposed to track those flow needs, and recommendations for additional research, monitoring, and analysis to improve understanding of flow needs. The report is part of the broader effort to identify, protect, and, where necessary, restore the Potomac River watershed's environmental flows.

Several typographical errors (pages 4, 5, 6, 53, 72, and 77) that appeared in the report published in May 2011 have been corrected in this version.

### Disc Contents

Document (PDF files)

- [AppendixA\\_LargeRiverEnvironFlowNeedsRpt.pdf](#) (100 KB)
- [Potomac Lg River Env Flows Final Combined\\_March-2012.pdf](#) (10,487 KB)

# Appendix B – Water Withdrawals and Consumptive Use in the Potomac River Basin Report

---

## Contents

This appendix contains the *Water Withdrawals and Consumptive Use in the Potomac River Basin* report. This stand-alone report describes how future water withdrawal and consumptive use for the year 2030 was estimated for six scenarios:

- high domestic and public water use (base scenario),
- medium domestic and public water use (DP2 scenario),
- low domestic and public water use (DP1 scenario),
- advanced technologies in the power sector and a new power generation facility (power scenario),
- conditions expected with climate change (climate change scenario), and
- drought conditions (hot and dry scenario).

The results were used to develop model inputs for future hydrologic alteration scenarios for other tasks in the Middle Potomac River Watershed Assessment project. Methodology for assembling datasets for those future flow scenarios is described in Appendix C, and methodology for how the future scenarios were modeled to produce flow time series is described in Appendix E. The study described here also serves as an update to a similar study done previously by ICPRB (Steiner et al. 2000). The two studies had different purposes and somewhat different datasets but some useful comparisons are made.

### Disc Contents

Document (PDF files)

- AppendixB\_FutureProjections.pdf (99 KB)
- FutureFlows\_FINAL.pdf (2,011 KB)

# Appendix C – Compilation of Measured Stream Data

---

## Contents

This appendix discusses in more detail sources of the data and GIS layers used in the project, and the analyses done with them. It also lists the data files provided in separate directories on the attached disc. Section names in the AppendixC\_Data.pdf document are identical to the directory names on the disc.

### Disc Contents

Document (PDF file)

- AppendixC\_Data.pdf (574 KB)

Eleven directories with data files and additional documentation (1.77 GB)

- 1\_Impoundments
- 2\_Withdrawals
- 3\_Discharges
- 4\_Land Cover
- 5\_Potomac-Susquehanna Flow Gages
- 6\_Stream Macroinvertebrates
- 7\_Model Input-Output
- 8\_Flow Metrics
- 9\_Future Scenarios
- 10\_GIS\_Files
- 11\_Master Data Set

# Appendix D – Application of ELOHA

---

## Contents

This appendix provides a more detailed description of the modification and application of the Ecological Limits of Hydrologic Alteration framework. This chapter was originally included as part of the main study report and followed section 5.1.

### Disc Contents

Document (PDF file)

- [AppendixD\\_Application\\_of\\_ELOHA.pdf](#) (1,819 KB)

# Appendix E – Development and Refinement of Hydrologic Model

---

## Contents

Hydrologic modeling efforts for the MPRWA project had multiple components. Detailed descriptions about the hydrologic modeling methodology are provided in the files on the attached disc. The file names listed below are identical to the file names on the disc.

### Disc Contents

Documents (PDF files)

- AppendixE\_HydrologicModel.pdf (74 KB)
- FutureScenarios\_011912.pdf (1,891 KB)  
Describes the development of the five future scenarios in the CBP HSPF modeling environment. It also spatially presents the resulting hydrologic alteration in seven selected flow metrics.
- Watershed\_delineation\_011712.pdf (537 KB)  
Describes how biological monitoring points were selected from the Chessie BIBI database and, once selected, how the watersheds draining to those locations were delineated.
- Resegmentation\_at\_Impoundments\_011712.pdf (376 KB)  
Describes the methodology utilized to re-segment the CBP HSPF model at “significant” impoundments in the study area.
- Pot-Susq\_CART\_analysis\_011712.pdf (486 KB)  
Describes how thresholds of flow alteration risk were identified by a Category and Regression Tree (CART) Analysis of the Potomac-Susquehanna dataset.
- Baseline\_Landuse\_011212.pdf (143 KB)  
Describes the calculation of land uses in the baseline model scenario.
- Baseline\_Scenario\_011212.pdf (580 KB)  
Describes the development of the baseline scenario in the CBP HSPF modeling environment including the removal of impoundments, withdrawals, and discharges, and conversion of current to baseline land uses. Select results of the baseline scenario are also presented.
- WOOOMM\_Inputs\_011012.pdf (525 KB)  
Describes the inputs needed to establish the ELOHA watersheds in the VADEQ WOOOMM environment and documents how those inputs were developed.
- Application\_of\_ModelingTools\_011312.pdf (619 KB)  
Describes the evaluation of the CBP Phase 5 model and VADEQ WOOOMM module for use in the Middle Potomac River Watershed Assessment.

# Appendix F – Flow Metric Testing

---

## Contents

This appendix provides a more detailed description of the analyses performed to verify the ability of the flow metrics generated from simulated time series to accurately represent those generated from observed time series.

### Disc Contents

Document (PDF file)

- [AppendixF\\_FlowMetricTesting.pdf](#) (1,819 KB)

# Appendix G – Stream Classification

---

## Contents

This appendix contains a report written by Julie Zimmerman (previous of The Nature Conservancy) in 2010 and titled *Susquebanna River stream classification analysis: determining a stream classification system to apply to the Potomac River and Chesapeake Bay*. The analysis described in the report was intended to help determine the need for stream classifications in the ELOHA analysis of the MPRWA project.

### Disc Contents

Document (PDF file)

- [AppendixG\\_StreamClassification.pdf](#) (189 KB)

# Appendix H – Basin-Wide Hydrologic Alteration Assessment

---

## Contents

The Middle Potomac River Watershed Assessment included a basin-wide assessment of hydrologic alteration. The purpose of this assessment was to understand the watershed drivers of hydrologic alteration (land uses, water uses, and meteorological conditions). The appendix describes in detail the methodology, sources, magnitude, and spatial extent of hydrologic alteration under current and future conditions in the study area, and the assessment results. It also provides an Excel file of the data and graphs used in the comparison of observed and simulated flow metrics plotted against different land uses.

### Disc Contents

Document (PDF file)

- Appendix H\_FlowAlterationAssessment.pdf (3,507 KB)

Data file (Excel 2007 spreadsheet)

- FlowAlt-vs-LandWater.xlsx (2,651 KB)

# Appendix I - Development of Stream Flow Alteration–Ecology Relationships

---

## Contents

This appendix provides detailed documentation on how the flow alteration – ecology (FA-E) relationships between the flow metrics and macroinvertebrate metrics were developed, and lists the files provided in separate directories on the attached disc.

### Disc Contents

Document (PDF file)

- AppendixI\_FA-E.pdf (234 KB)
- FA-E\_ CondProb\_plots.pdf (398 KB)
- FA-E\_QuantileRegr\_plots.pdf (433 KB)

Data file (Excel 2007 spreadsheet)

- QuantileRegr\_Data\_R-scripts.xlsx (27 KB)
- CondProb\_Data\_R-scripts.xlsx (1,412 KB)
- PearsonCorrelations.xlsx (1,628 KB)

# Appendix J - Stakeholder Engagement

---

## Contents

One objective of this project was a transfer of knowledge gained to “stakeholders, other interested parties, and USACE Districts and Divisions”. In addition, other tasks included elements specifying consultation and/or review of interim products by stakeholders or experts. The project team addressed these requirements by building a website for sharing information about the project, presenting webinars, holding two workshops, and establishing a Technical Advisory Committee and having two meetings with that group.

### Disc Contents

Document (PDF file)

- AppendixJ\_StakeholderEngagement.pdf (146 KB)
- Potomac PMP signed.pdf (815 KB)
- MPRWA\_fact\_sheet\_2011.pdf (591 KB)
- MPRWA\_Intro\_09\_29\_09.pdf (4,470 KB)
- LgRivWorkshopIntro\_090910.pdf (3,841 KB)
- WebinarSeriesIntro\_030811.pdf (5,667 KB)
- TechOverview\_No1\_041211\_V3.pdf (1,110 KB)
- HumanUsesOfWater\_No2\_051011.pdf (2,073 KB)
- ModelingStreamflow\_No3\_061611.pdf (2,484 KB)
- FlowEcology1\_No4\_071411.pdf (3,166 KB)
- FlowEcology2\_No5\_090811.pdf (3,874 KB)
- Science\_to\_Mgmt\_No6\_102711.pdf (3,175 KB)
- FutureScenarios\_No7\_022312.pdf (2,341 KB)
- FinalWebinar\_No8\_06-21-12.pdf (2,066 KB)
- MPRWA\_ATR.pdf (163 KB)
- MPRWA\_constituent\_comments\_10\_12\_2012.pdf (175 KB)

Three directories (28.7 MB)

- TAG - contains materials presented at 28 September and 26 October, 2011 meetings of the Technical Advisory Group
- Workshop – contains materials presented at the 29-30 November, 2011 workshop
- Correspondence – contains copies of correspondence to federal and state agencies regarding the Middle Potomac River Watershed Assessment

# Appendix K – Implementation Options

---

## Contents

This appendix documents implementation efforts that have been investigated as part of the Middle Potomac River Watershed Assessment, including a literature review of Decision Support Tools (DST\_CaseStudies.pdf) and a concept paper that scopes out a basin-wide comprehensive plan for the Potomac Basin (CompPlan\_ConceptPaper\_042911.pdf).

### Disc Contents

Document (PDF file)

- AppendixK\_Implementation.pdf (69 KB)
- DST\_CaseStudies.pdf (6,190 KB)

One implementation consideration that underwent evaluation was the development of a computer-based decision making tool. To this end, case studies of Decision Support Tools from across the country were compiled and are contained in this document.
- CompPlan\_ConceptPaper\_042911.pdf (121 KB)

The scoping phase of a Potomac Basin Comprehensive Water Resource Plan was completed as a part of the Middle Potomac River Watershed Assessment. This document outlines the planning process, as currently scoped. A version of this document was published in the AWRA 2011 summer specialty conference proceedings, held in Snowbird, Utah on June 27-29, 2011.